

Vaccine Clinic Resource for Immunizers

Disclaimer: this Quick Reference is not intended to replace other product specific vaccine references. The document is intended as a quick reference for frequently referred to information. Please refer to the product monograph for all current and complete information.

T'11 -	
Title:	Flu & Pneumo Vaccine Quick Reference Guide 2023-2024
Flu vaccine	A/Victoria/4897/2022 (H1N1)pdm09-like virus
compositions comply	A/Darwin/9/2021 (H3N2)–like virus
with the WHO's	B/Austria/1359417/2021 (B/Victoria lineage)-like virus
recommendations for	B/Phuket/3073/2013 (B/Yamagata lineage)-like virus
the 2023-2024 season:	
Formulations:	IIV4-SD (Inactivated Influenza Vaccine Quadrivalent- Standard dose)
	IIV4-HD (Inactivated Influenza Vaccine Quadrivalent- High dose)
	Pneu-P-23 (Penumococcal Polysaccharide)
Eligibility Criteria:	Seasonal Flu Health Province of Manitoba (gov.mb.ca)
	Vaccine Eligibility Health Province of Manitoba
Fact Sheets:	Seasonal Influenza Vaccine
	High Dose Seasonal Influenza Vaccine for People Aged 65 years and Older
	Polysaccharide (Pneu-P-23) Vaccine
Canadian Immunization	Influenza Vaccine: Influenza vaccine: Canadian Immunization Guide - Canada.ca
Guide:	Pneumococcal Vaccine: Pneumococcal vaccine: Canadian Immunization Guide - Canada.ca

Note: Egg allergic individuals may be vaccinated using any of the seasonal influenza vaccines available in MB, without prior influenza vaccine skin test and with the full dose, irrespective of a past severe reaction to egg, and without any extraordinary precautions, but ensuring that immunizers be prepared with the necessary equipment, knowledge and skills to respond to a vaccine emergency.



Presentation	Storage and Handling	Administration	Ingredients/
		(Dosage/route/schedule)	Potential Allergens
Supplied: 0.5ml single dose prefilled syringes (package of 10) OR 5 ml multidose vial (10 doses) Shake well to uniformly distribute the suspension and inspect visually before administering each dose. After shaking well, solution is clear to slightly opalescent in colour. There should NOT be any particulate matter and/or discolouration- if these exist, do not administer. Product Monograph: https://www.gov.mb.ca/health/flu/docs/pm/fluzone-qiv.pdf	Storage: 2 to 8 degrees C° Do not freeze Discard product if exposed to freezing Once punctured, the multi-dose vial can be used until expiry date. Unpunctured vial is stable until expiry date. A maximum of 10 doses can be withdrawn from a multidose vial Protect from light	6 months to 8 years of age: Dose: 0.5ml dose Route: → 6m to less than 12m of age: IM Vastus lateralis → 12m and older: IM Deltoid Schedule: 1 or 2 doses Children 6m to 8 years of age who have not previously been vaccinated against influenza should receive a second dose at least 4 weeks later Recommended Needle Gauge and Length: 22-25G 5/8"-1" 9 years of age and older: Dose: 0.5ml dose Route: IM Deltoid Schedule: 1 dose Recommended Needle Gauge and Length: 22-25G 1"-1 ½"	Egg proteins Formaldehyde Sodium phosphate-buffered Isotonic sodium chloride Triton® X-100 Thimerosal (in the multidose vial only) LATEX FREE



Presentation	Storage and Handling	Administration (Dosage/route/schedule)	Ingredients/ Potential Allergens
Supplied: 5 ml multidose vial (10 doses) Shake well prior to each administration and inspect visually for any foreign particulate matter and/or variation of physical aspect prior to administration. In the event of either, discard the vaccine. Liquid should be opalescent translucent to off-white suspension that may sediment slightly. Product Monograph: https://www.gov.mb.ca/health/flu/docs/pm/flulaval-tetra.pdf	Storage: 2 to 8 degrees C° Do not freeze Discard product if exposed to freezing The vaccine is stable for 12 months. Do not use past the expiry date. Once punctured, the multi-dose vial should be discarded within 28 days. Protect from light	6 months to 8 years of age: Dose: 0.5ml dose Route: → 6m to less than 12m of age: IM Vastus lateralis → 12m and older: IM Deltoid Schedule: 1 or 2 doses Children 6m to age 8 who have not previously been vaccinated against influenza should receive a second dose at least 4 weeks later Recommended Needle Gauge and Length: 23-25G 5/8"-1" 9 years of age and older: Dose: 0.5ml dose Route: IM Deltoid Schedule: 1 dose Recommended Needle Gauge and Length: 23-25G 1"-1 ½"	 α- tocopheryl hydrogen succinate Polysorbate 80 Phosphate buffered saline Thimerosal Trace amounts of: Egg proteins Ethanol Formaldehyde Sodium deoxycholate Sucrose LATEX FREE



Presentation	Storage and Handling	Administration (Dosage/route/schedule)	Ingredients/ Potential Allergens
Supplied: 0.5ml single dose prefilled syringes (package of 10) OR 5 ml multidose vial (10 doses) Shake well to uniformly distribute the suspension and inspect visually before administering each dose. The vaccine is a slightly opaque liquid with some white particulate sediment that resuspends upon shaking to form a homogenous suspension. Do not use if extraneous particulate matter and/or discolouration is observed. Product Monograph: https://www.gov.mb.ca/health/flu/docs/pm/afluria-tetra.pdf	Storage: 2 to 8 degrees C° Do not freeze Discard product if exposed to freezing The shelf-life of the vaccine is 12 months. Do not use past the expiry date. Unpunctured vial is stable until expiry date on package. Once punctured, the multidose vial must be used within 28 days. A maximum of 10 doses can be withdrawn from a multidose vial (do not exceed 10 needle punctures) Protect from light	5 to 8 years of age: Dose: 0.5ml dose Route: IM Deltoid Schedule: 1 or 2 doses Children age 5 to age 8 who have not previously been vaccinated against influenza should receive a second dose at least 4 weeks later Recommended Needle Gauge and Length: 22-25G 7/8"-1" 9 years of age and older: Dose: 0.5ml dose Route: IM Deltoid Schedule: 1 dose Recommended Needle Gauge and Length: 22-25G 1"-1 ½"	Calcium chloride Dibasic sodium phosphate Monobasic sodium phosphate Monobasic potassium phosphate Potassium chloride Sodium chloride Thimerosal (multidose vial only) May also contain trace amounts of: Sodium Taurodeoxycholate Beta-propiolactone Hydrocortisone Sucrose Neomycin Sulfate Polymyxin B Sulfate Egg Protein LATEX FREE



Fluzone® High-Dose Quadrivalent (IIV4-HD): Use in 65 years of age and older Sanofi Pasteur

Presentation	Storage and Handling	Administration (Dosage/route/schedule)	Ingredients/ Potential Allergens
Supplied: 0.7ml single dose prefilled syringes (package of 10)	Storage: 2 to 8 degrees C°	65 years of age and older: Dose: 0.7ml dose	 Triton® X-100 Sodium phosphate-buffered isotonic sodium chloride solution
Shake well to uniformly distribute the suspension and inspect visually before administering each dose. After shaking the syringe, the liquid should be clear and slightly opalescent in colour.	Do not freeze Discard product if exposed to freezing Do not use after the	Route: IM Deltoid Schedule: 1 dose	Trace amounts of: Egg Protein Formaldehyde
Do not use if extraneous particulate matter and/or discolouration is observed.	expiration date shown on the label	Recommended Needle Gauge and Length: 22-25G 1"-1 1/2"	LATEX FREE
Product Monograph: https://www.gov.mb.ca/health/flu/docs/pm/fluzone-hd-qiv.pdf	Protect from light		

Pneumovax 23® (Pneu-P-23 Penumococcal Polysaccharide): Use in 65 years of age and older And those 2 years of age and older who meet the <u>Manitoba Health Eligibility Criteria</u>

Merck Canada

Presentation	Storage and Handling	Administration (Dosage/route/schedule)	Ingredients/ Potential Allergens
Supplied: 0.5ml single dose prefilled syringes	Storage: 2 to 8 degrees C°	2 years of age and older:	Sodium chloride Phenol
Product Monograph:		Dose: 0.5ml	
[Product Monograph Template - Schedule D] (hres.ca)	Do not freeze Discard product if exposed to freezing	Route: IM Deltoid OR Subcutaneous (SC) Schedule: 1 dose	LATEX FREE
	Do not use after the expiration date shown on the label	A second dose may be recommended for those at highest risk. This should be assessed by the primary health care provider.	
	Protect from light		
		Recommended Needle Gauge and Length: IM: 22-25G 1"-1 1/2" SC: 25G 5/8"	