

Manitoba Health STI Medication Order Form (PLEASE PRINT)



Fax order to: Taché Pharmacy – Fax: (204) 231-1739 • Phone: (204) 233-3469 • For Inquiries only e-mail tache@mts.net

| Date (yyyy/mm/dd) | ____/____/____ | Contact Person: | |
|--|---|--|--|
| Facility Name: | | Physician: | |
| Address: | | | |
| Telephone #: | | Ext: | |
| Refer to Sexually Transmitted and Blood-Borne Infections webpage https://www.gov.mb.ca/health/publichealth/cdc/sti/index.html | | | |
| All treatments provided to patients must be recorded and reported to Manitoba Health using the Provider Report Form for Sexually Transmitted and Blood-Borne Infections (STBBI) and STI Treatment (https://www.gov.mb.ca/health/publichealth/surveillance/docs/mhsu_6781.pdf) | | | |
| Units Ordered | Drug | Dosage Forms & Unit Quantity | Indication |
| CHANCROID | | | |
| | Ceftriaxone | 250 mg x 1 vial | First line treatment for adults and adolescents |
| | Erythromycin base | 250 mg x 56 tablets | Alternative treatment for cephalosporin or penicillin allergy. |
| | Azithromycin | 250mg x 4 tablets | |
| CHLAMYDIA[†] | | | |
| | Azithromycin | 250 mg x 4 tablets | First line treatment in children > 9 years of age and adults |
| | Doxycycline | 100 mg x 14 capsules | Alternative treatment in children > 9 years of age and adults |
| | Erythromycin base | 250 mg x 56 tablets | Pregnant and lactating women |
| | Amoxicillin | 500 mg x 21 capsules | Pregnant and lactating women allergic to erythromycin or azithromycin |
| GONORRHEA[†] | | | |
| | Cefixime | 400 mg x 2 tablet | First line treatment in combination with 1 g azithromycin |
| | Ceftriaxone | 250 mg x 1 vial | First line treatment in combination with 1 g azithromycin |
| | Azithromycin | 250 mg x 4 tablets | First line treatment treatment in combination with cefixime OR ceftriaxone |
| | Azithromycin | 250 mg x 8 tablets | Alternative treatment when there is severe cephalosporin allergy |
| LGV | | | |
| | Doxycycline | 100 mg x 42 capsules | First line treatment |
| | Erythromycin base | 250 mg x 168 tablets | Alternative treatment |
| | Azithromycin | 250 mg x 12 tablets | Alternative treatment |
| PID | | | |
| | Ceftriaxone | 250 mg x 1 vial | Recommended treatment for acute PID (in combination with doxycycline and metronidazole) |
| | Doxycycline | 100 mg x 28 capsules | Recommended treatment for acute PID (in combination with ceftriaxone and metronidazole) |
| | Metronidazole | 250 mg x 56 tablets | Recommended treatment for acute PID (in combination with doxycycline and ceftriaxone). Metronidazole is included to provide optimal anaerobic coverage. |
| | Azithromycin | 250 mg x 7 tablets (1 tablet daily for 7 days) OR | Alternative treatment for acute PID (in combination with oral metronidazole) for patients with cephalosporin contraindication. |
| | Azithromycin | 250 mg x 8 tablets (4 tablets once weekly for 2 weeks) | |
| SYPHILIS | | | |
| | Benzathine Penicillin G [†] (Bicillin) | 1.2 MU x 2 syringes | First line for primary, secondary, and early latent stages of infection and epidemiologic treatment of sexual contacts. |
| | Benzathine Penicillin G [†] (Bicillin) | 1.2 MU x 6 syringes | First line for late latent infection, infection of unknown duration and tertiary syphilis NOT involving the CNS. |
| | Ceftriaxone | 1 g x 10 vials | Alternative treatment for penicillin allergy. |
| | Doxycycline | 100 mg x 28 capsules | Alternative treatment for penicillin allergic adults and adolescents staged with primary, secondary or early latent infection |
| | Doxycycline | 100 mg x 56 capsules | Alternative treatment for penicillin allergic adults and adolescents staged with late latent infection, infection of unknown duration or tertiary syphilis NOT involving the CNS |
| ^{††} Pediatric dosing available upon request. [†] Must be stored in a refrigerator that is temperature monitored and kept between 2-8°C. | | | |
| Prescriber's Name: | | Prescriber's License No: | Prescriber's Signature: |