- May 19-25: The proportion of patients visiting sentinel physicians for influenza-likeillness was 5.1% (up from 2.0% last week).
- •This week: There were **0** cases of influenza A and **1** case of influenza B reported.
- •A total of **590** cases of influenza A and **111** cases of influenza B have been reported since the start of the current influenza season.
- There was 1 hospitalization associated with a laboratory-confirmed diagnosis of influenza that were reported this week, 1 ICU admission and 0 deaths.
- There have been 152
 hospitalizations, of which 25
 resulted in ICU admissions, this
 season.
- So far this season, 9
 Manitobans with laboratory-confirmed influenza have died.

Severity



Outpatient ILI (sentinels)



Laboratory

- •The following regional proportions are observed among cases of influenza A: Winnipeg (34.7%), Northern (13.9%), Prairie Mountain (26.4%), Southern (16.0%), Interlake-Eastern (9.0%).
- Between Jan 1 May 26, 2013, 1336 units of oseltamivir have been dispensed from community retail pharmacies. This brings the total number of units dispensed since Sept 1, 2012 up to 1573.
- •There has been 1 H3N2 influenza isolate that has tested positive for resistance to Oseltamivir and Zanamivir and 1 H1N1 influenza isolate that tested positive for resistance to Oseltamivir.

Geography



Treatment



Antiviral Resistance



- Manitoba's influenza activity, as estimated by Google search data, has been downgraded to "minimal".
- •There were 8 calls to Health Links - Info Santé this week, which is about the same as the previous week (10 calls).

Syndromic Surveillance



 As of May 25, there have been
 40 lab-confirmed outbreaks of influenza A reported, of which one is ongoing.

Institutional
Outbreaks



•As of April 6, 2013, only **20.1%** of Manitobans had received the seasonal influenza vaccine.

Immunization







In Summary

• There were no laboratory-confirmed cases of influenza A and 1 case of influenza B reported last week.

Surveillance Measures

1. Laboratory Surveillance

Reports of culture isolations and enzyme immunoassay (EIA) detections from Cadham Provincial Laboratory (CPL) are forwarded to the Public Health Surveillance (PHS) Unit weekly. While EIA detections and culture isolations comprise the largest number of reports from CPL, seroconversions are similarly forwarded to the PHS Unit weekly.

Age Group, Maintoba, 2012/2015							
Age	Influ	enza A	Influenza B				
Group	#	%	#	%			
<1 yrs	42	7.1%	5	4.5%			
1-4 yrs	61	10.3%	8	7.2%			
5-9 yrs	40	6.8%	11	9.9%			
10-14 yrs	23	3.9%	10	9.0%			
15-19 yrs	32	5.4%	6	5.4%			
20-24 yrs	22	3.7%	4	3.6%			

3.2%

4.6%

7.1%

8.5%

6.6%

8.3%

24.4%

0.0%

6

9

9

14

8

4

17

111

5.4%

8.1%

8.1%

12.6%

7.2%

3.6%

15.3%

0.0%

19

27

42

50

39

49

144

590

25-29 yrs

30-39 yrs

40-49 yrs

50-59 yrs

60-69 yrs

70-79 yrs

>79 yrs

Missing

TOTAL

Table 1. Reported Cases of Influenza A and B by

This week, there were:

- 0 cases of influenza A reported;
- 1 case of influenza B reported.

Since the beginning of this season, there have been:

- 590 cases of influenza A reported;
- 111 cases of influenza B reported.

May 19-25, 2013

Figure 3. Number of lab-confirmed cases of influenza A by specimen collection week and season, Manitoba (*Note:* 2009/2010 season excluded due to the H1N1 pandemic, making the numbers incomparable with other seasons.)





Figure 4. Number of lab-confirmed cases of influenza B by specimen collection week and season, Manitoba








2. Outpatient ILI (Sentinel Physicians)

Manitoba Health participates in the National *FluWatch* Program coordinated by PHAC. In addition to laboratory-confirmation of influenza, this program relies on weekly reports of ILI as reported by 22 current Manitoban sentinel physicians in all five RHAs (Northern, Southern, Prairie Mountain, Interlake-Eastern, and Winnipeg).

The proportion of patients seen for an ILI this week was higher than last week (5.1% from 2.0%). The proportion is higher than what was observed at the same time last season.

Manitoba Health receives weekly reports from PHAC presenting the provincial ILI rate and the specific data for each of the participating sentinel physicians. The graph below depicts the proportion of total patients who were seen for an ILI. These numbers should be interpreted with caution as the number of sentinel physicians reporting to *FluWatch* varies from week to week and may not be representative of ILI activity across the province.

Figure 1. Proportion of patients seen for ILI as reported by *FluWatch* sentinel physicians by week and flu season, Manitoba



STRIVE (Surveillance Team Research on Influenza Vaccine Effectiveness)

Beginning with the 2012/2013 influenza season, Manitoba Health has joined STRIVE, a national multisite vaccine effectiveness surveillance network already in operation in Alberta, British Columbia, Ontario, and Quebec. Operated in collaboration with Cadham Provincial Laboratory, STRIVE aims to assess the effectiveness of the 2012-2013 seasonal trivalent influenza vaccine in protecting against influenza, and to monitor influenza activity in the region. STRIVE specimens are tested for influenza and other respiratory viruses through PCR and Seeplex RV15 panel. Results of respiratory testing performed by network members will be regularly featured in this column. While recruitment is ongoing, we would like to thank sentinel clinicians and sites who have thus far supported this public health initiative in Manitoba. For more information about the study, please e-mail Arielle.GoldmanSmith@gov.mb.ca (for sites outside Winnipeg) or strive@wrha.mb.ca (Winnipeg).

	Influenza A	Influenza B	Total			
*New STRIVE lab-confirmed influenza cases:	0	1	1			
Total STRIVE lab-confirmed influenza cases: 46 13 59						
(These cases are included in the total number of provincial influenza cases)						


4

Health Links – Info Santé (HL-IS)

HL-IS is a 24-hour, 7-days a week telephone information service staffed by registered nurses with the knowledge to provide answers over the phone to health care questions and guidance to appropriate care. When a caller phones HL-IS and selects the Influenza Service, they are given an option to select information on (1) the groups of individuals who are at an increased risk of serious illness, (2) how to arrange a flu shot, (3) the annual influenza immunization campaign, or (4) the management of flu and its potential complications.

This week there were 8 calls, which is about the same as the previous week (10 calls). The weekly total is similar to the total observed at the same time last season.

Figure 2. The number of calls to HL-IS Influenza Service by season, Manitoba



Severity (Clinically Severe Cases)

This season, PHAC is requesting provinces and territories to report the number of hospitalizations, ICU admissions, and deaths <u>associated</u> with a lab-confirmed report of influenza. The reason for hospitalization, ICU admission, or death does not have to be attributable to the influenza diagnosis in order to be included in this count. These data are collected in order to continue with the surveillance system implemented during the 2009 H1N1 pandemic to help monitor the severity/burden of illness during the influenza season.

This week there were:

- *Since the beginning of the season, there have been:
- 1 hospitalization, of which
- 1 resulted in an ICU admission; and
- 0 deaths.¹

- 152 hospitalizations, of which
- 25 resulted in an ICU admission; and
- 9 deaths.¹

There were 47 children (aged 9 or under) admitted to hospital with laboratory-confirmed influenza A or B since the start of the season. There was one child admitted in the week of May 19-25, 2013. (Note: These children are included in the counts above).

*Hospitalized cases are reported based on laboratory report date.

¹ The reason for hospitalization, ICU admission, or death does not have to be attributable to the influenza diagnosis in order to be included in this count.

Institutional Outbreaks

Outbreaks of influenza must be accompanied by a positive influenza lab report to be counted. The outbreak-related cases reflected on tables and figures within this report are lab-confirmed. However, most outbreak-related cases will not be lab-confirmed.

This week there were:

From the beginning of the season until May 25, there have been:

- 0 outbreaks of influenza A;
- 0 outbreaks of influenza B.

- 40 outbreaks of influenza A;
- 5 outbreaks of influenza B.


Table 2. Cumulative number of lab-confirmed outbreaks of influenza by RHA and season, Manitoba

	2011/2012 up to the end of Week 21 (May 26, 2012) 2012/2013 up to end of the				the Week 2	21 (May 25, 2013)		
RHA:	LTCF	Workplace	ACF	Community	LTCF	Workplace	ACF	Community
Winnipeg	8	0	0	0	23	0	0	0
Northern	0	0	0	0	2	0	0	0
Southern	0	0	0	0	5	0	0	0
Interlake-Eastern	1	0	0	0	4	0	0	0
Prairie Mountain	0	0	0	0	11	0	0	0
Total	9	0	0	0	45	0	0	0

LTCF: long term care facility ACF: acute care facility

Syndromic Surveillance

<u>Google Flu Trends</u> uses aggregated Google search data to estimate influenza activity. As of May 30, Manitoba's influenza activity has been downgraded from "low" to "minimal".



Sub-Typing, Strain Characterization, and Antiviral Resistance

Sub-Typing:

Table 3. Sub-typing of influenza A specimens as reported by CPL, 2012/2013 flu season, Manitoba

A/H1	A/H3	A Unsubtyped	A Total
18	167	405	590

Strain Characterization:

Since September 1, 2012, NML has antigenically characterized 1309 influenza viruses (630 H3N2, 206 H1N1, and 473 B viruses) that were received from Canadian laboratories with the following results:

Strain	Number of viruses		
	Canada	Manitoba	
A/Victoria/361/2011 (H3N2)-like ²	630	3	
A/California/07/09 (H1N1)-like ³	206	0	
B/Brisbane/60/2008-like (B/Victoria/02/87 lineage) ⁴	103	2	
B/Wisconsin/01/2010-like (Yamagata lineage) ⁵	370	4	

Antiviral Resistance:

Since September 1, 2012, NML has performed drug susceptibility testing on influenza isolates received from Canadian laboratories with the following results:

Antiviral resistance by influenza virus type and subtype, Canada, 2012/2013							
Virus	Oseltamivir		Zanamivir		Amantadine		
type/subtype	# Resistant (%)	# Sensitive (%) # Resistant (%) # Sensitive (%)		# Resistant (%)	# Sensitive (%)		
A(H3N2)	1 (0.2)	623 (99.8)	1 (0.2)	618 (99.8)	1011 (99.9)	1 (0.1)	
A(H1N1)	1 (0.5)	206 (99.5)	0	200 (100)	244 (100)	0	
В	0	468 (100)	0	468 (100)	N/A	N/A	

N/A = Not applicable

The isolates tested from CPL had the following results:

Antiviral resistance by influenza virus type and subtype, Manitoba, 2012/2013							
Virus	Oseltamivir		Zanamivir		Amantadine		
type/subtype	# Resistant	# Sensitive	# Resistant # Sensitive		# Resistant	# Sensitive	
A(H3N2)	0	3	0	3	6	0	
A(H1N1)	0	0	0	0	2	0	
В	0	4	0	4	N/A	N/A	

² Strain match to recommended H3N2 component for the 2012/2013 northern hemisphere influenza vaccine.

³ Strain match to recommended H1N1 component for the 2012/2013 northern hemisphere influenza vaccine.

⁴ Strain match to recommended influenza B component of the 2011/2012 influenza vaccine.

⁵ Strain match to recommended influenza B component for the 2012/2013 northern hemisphere influenza vaccine.

Abbreviations

ACF = acute care facility

CPL = Cadham Provincial Laboratory

HL-IS = Health Links - Info Santé

PHAC = Public Health Agency of Canada

ICU = intensive care unit

ILI = influenza-like-illness

LTCF = long term care facility

NML = National Microbiology Laboratory

PHS = Public Health Surveillance

RHA = Regional Health Authority

WRHA = Winnipeg Regional Health Authority

Explanatory Notes and Definitions

Cumulative data:

Cumulative data includes updates to previous weeks; due to reporting delays or amendments, the sum of weekly report totals may not add up to cumulative totals.

Data extraction date:

Manitoba-specific information contained within this update is based on data confirmed in Manitoba's PHS Unit databases on or before **May 31, 2013**, the date of data extraction.

ILI in the general population:

Acute onset of respiratory illness with fever and cough and with one or more of the following – sore throat, arthralgia, myalgia, or prostration, which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

ILI outbreaks:

Schools: Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI.

Hospitals and residential institutions: Two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case.

Other settings: Two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. workplace, closed communities.

Specimen collection date:

The date the laboratory specimen is taken is used to assign cases to the appropriate week in this report. However, hospitalized/ICU cases are reported based on laboratory report date.

For other Epidemiology and Surveillance reports, please view the Manitoba Health internet website: http://www.gov.mb.ca/health/publichealth/surveillance/index.html

For national surveillance data, refer to:

http://www.phac-aspc.gc.ca/fluwatch/index-eng.php