



Winnipeg Regional Health Authority  
*Caring for Health*

Office régional de la santé de Winnipeg  
*À l'écoute de notre santé*

490 Hargrave Street  
Winnipeg, Manitoba  
R3A 0X7 CANADA

[www.wrha.mb.ca](http://www.wrha.mb.ca)

490, rue Hargrave  
Winnipeg (Manitoba)  
R3A 0X7 CANADA

[www.wrha.mb.ca](http://www.wrha.mb.ca)

November 26, 2020

John Taylor Collegiate  
470 Hamilton Ave.  
Winnipeg, MB  
R2Y 0H4

Dear Parent/Guardian of Students at John Taylor Collegiate,

Manitoba public health officials advised John Taylor Collegiate today of a confirmed case of COVID-19 in the school on November 19 when the individual may have been infectious.

John Taylor Collegiate is working closely with public health officials and following their recommendations. Based on the public health investigation, this exposure was assessed to be low risk. The infection was not believed to be acquired at school. At this time, no close contacts have been identified at John Taylor Collegiate. No one is required to self-isolate from this exposure.

Please note: the school is not permitted to release the name or any identifying information about the confirmed case of COVID-19 to any member of the school, community or media.

If your child develops a new onset or worsening of any one symptom listed in Column A or any two or more symptoms listed in Column B (see table below), even if they are mild, immediately isolate them from others. It is recommended your child get tested for COVID-19 as soon as symptoms appear. You may go directly to the nearest COVID-19 testing location during their open hours. For the testing site nearest you, please visit <https://www.gov.mb.ca/covid19/updates/testing.html>, or call Health Links—Info Santé at 204-788-8200 or toll-free at 1-888-315-9257.

A	B
Fever/Chills	Runny nose
Cough	Muscle aches
Sore throat/hoarse voice	Fatigue
Shortness of breath/difficulty breathing	Pink eye (conjunctivitis)
Loss of taste or smell	Headache
Vomiting or diarrhea for more than 24 hours	Skin rash of unknown cause
	Nausea or loss of appetite
	Poor feeding (if an infant)

Please note: individuals with symptoms outlined in columns A and B, who choose not to be tested, must isolate for a minimum of 10 days following the start of symptoms. Isolation can be stopped at day 10 if the individual has been symptom free for 24 hours.

For more public health information about COVID-19, please visit <https://manitoba.ca/covid19/updates/resources.html>.

Sincerely,



Dr. Heejune Chang  
 Medical Officer of Health  
 Winnipeg Regional Health Authority