



Winnipeg Regional Health Authority
Caring for Health

Office régional de la santé de Winnipeg
À l'écoute de notre santé

490 Hargrave Street
Winnipeg, Manitoba
R3A 0X7 CANADA

490, rue Hargrave
Winnipeg (Manitoba)
R3A 0X7 CANADA

www.wrha.mb.ca

www.wrha.mb.ca

November 9, 2020

Strathcona School
233 McKenzie St.
Winnipeg, MB
R2W 4Z2

Dear Parent/Guardian of students at Strathcona School,

Manitoba public health officials advised Strathcona School today of a confirmed case of COVID-19 present in the school on November 3 and 4, 2020 when the individual may have been infectious.

Strathcona School is working closely with public health officials and following their recommendations. Based on the public health investigation, this exposure was assessed to be low risk overall for the school community.

Close contacts of the cases have been identified and advised to self-isolate (quarantine). Anyone who was in the same room has been notified and is recommended to self-monitor for symptoms until November 18 and seek testing if symptoms develop.

Please note: the school is not permitted to release the name or any identifying information about the confirmed case of COVID-19 to any member of the school, community or media.

If your child develops a new onset or worsening of any one symptom listed in Column A or any two or more symptoms listed in Column B (see table below), even if they are mild, immediately isolate them from others. It is recommended your child get tested for COVID-19 as soon as symptoms appear. You may go directly to the nearest COVID-19 testing location during their open hours. For the testing site nearest you, please visit gov.mb.ca/covid19/locations.html, or call Health Links—Info Santé at 204-788-8200 or toll-free at 1-888-315-9257.

A	B
Fever/Chills	Runny nose
Cough	Muscle aches
Sore throat/hoarse voice	Fatigue
Shortness of breath/difficulty breathing	Pink eye (conjunctivitis)
Loss of taste or smell	Headache
Vomiting or diarrhea for more than 24 hours	Skin rash of unknown cause
	Nausea or loss of appetite
	Poor feeding (if an infant)

Please note: individuals with symptoms outlined in columns A and B, who choose not to be tested, must isolate for a minimum of 10 days following the start of symptoms. Isolation can be stopped at day 10 if the individual has been symptom free for 24 hours.

For more public health information about COVID-19, please visit <https://manitoba.ca/covid19/updates/resources.html>.

Sincerely,



Dr. Lisa Richards MD MSc FRCPC
 Medical Officer of Health
 Winnipeg Regional Health Authority