

Influenza Surveillance Report 2017–2018

Week 52 (Dec. 24–30, 2017)

Data extracted Jan. 5, 2018 at 11:00 am

Summary: High Activity

The seasonal influenza activity level in Manitoba is high. Influenza detections increased quickly across the whole province in the last two weeks. Influenza A and B are co-circulating and in Week 52, they together accounted for almost 80% of laboratory detections in respiratory viruses. There were 15 respiratory outbreaks reported in Week 52 and nine were confirmed to be influenza outbreaks. Two of these outbreaks were due to both influenza A and B. The Emergency Departments this week continued to have an increased number of patients with respiratory symptoms, especially at the Children's Hospital. In total, there have been 95 influenza associated hospital admissions since the beginning of this season and 65% of those patients are over the age of 65 years.

Influenza activity continues to increase across Canada overall. In many areas, the increase is consistent with the expected seasonal trend. The majority of influenza detections across Canada continue to be A(H3N2). A number of influenza B detections have also been reported to be substantially greater than in previous seasons. There have also been a small number of influenza A(H1N1) detections.

As of Dec. 10, 2017, influenza activity continued to increase in the temperate zone of the northern hemisphere, while in the temperate zone of the southern hemisphere, activity decreased at inter-seasonal levels according to the World Health Organization. In Central America and the Caribbean, influenza activity remained low. Worldwide, influenza A(H3N2) and B viruses accounted for the majority of influenza detections.

Laboratory

Laboratory-confirmed influenza cases this week:

- Influenza A cases: **147** ↑
- Influenza B cases: **10** ↑

Since Sept. 1, 2017:

- Influenza A cases: **381**
- Influenza B cases: **20**

Severity

Severe outcomes associated with laboratory-confirmed diagnosis of influenza this week:

- Hospitalizations: **18**
- ICU* admissions: **2**
- Deaths: **0**

Since Sept. 1, 2017:

- Hospitalizations: **95**
- ICU* admissions: **10**
- Deaths: **9**

Outbreak

Laboratory-confirmed Influenza outbreaks this week:

- Influenza A outbreaks: **7** ↑
- Influenza B outbreaks: **0**
- Influenza A and B outbreaks: **2** ↑

Since Sept. 1, 2017:

- Influenza A outbreaks: **23**
- Influenza B outbreaks: **0**
- Influenza A and B outbreaks: **2**

Syndromic in Community

Calls to Influenza Service at Health Links–Info Santé this week: **30**

Syndromic in Care

Visits to sentinel physicians due to ILI this week: **9.4%** ↑

Units of antiviral dispensed from retail pharmacies this week: **108** ↑

Syndromic in ED

Respiratory visits to Emergency Department (ED) this week: **214/day** ↑

Immunization

As of Dec. 29, 2017:

- Percentage of Manitoban residents immunized with the seasonal influenza vaccine: **19.6%**

Antiviral Susceptibility

Isolates resistant to antiviral since Sept. 1, 2017 in Canada:

- Oseltamivir: **0** (out of 287 tests)
- Zanamivir: **0** (out of 226 tests)

Regional Health Authority

Influenza cases (cases/100,000 population) since Sept. 1, 2017:

- Winnipeg: **194 (25.3)**
- Southern: **33 (16.7)**
- Interlake-Eastern: **29 (22.6)**
- Prairie Mountain: **88 (51.8)**
- Northern: **57 (74.6)**

Note. * ICU admissions were also included in hospitalizations.

Time trends in this report were analyzed by [epidemiology week](#).

Numbers are subject to change. Missed events in the current report due to a delay of submission to MHSAL will be included in later reports when data become available.

Figure 1. Weekly Cases of Laboratory-Confirmed Influenza, Manitoba




Figure 2. Weekly Influenza and ILI Outbreaks, Manitoba




Figure 3. Weekly Influenza Related Calls to Health Link – Info Santé, Manitoba




Figure 4. Weekly % of ILI Related Visits to Sentinel Physicians, Manitoba




Figure 5. Weekly Units of Antiviral Dispersed from Pharmacies, Manitoba




Figure 6. Weekly Respiratory Visits to Emergency Department and % of Total, Manitoba, 2017-2018




Figure 7. Weekly Total and ILI Visits to Emergency Department, Manitoba




Table 1. Antiviral Resistance of Isolates by Influenza Type and Subtype since September 1, 2017, 2017–2018

		Oseltamivir		Zanamivir		Amantadine	
		Resistant	Sensitive	Resistant	Sensitive	Resistant	Sensitive
Canada	A(H3N2)	0	206	0	168	173	0
	A(H1N1)	0	20	0	12	12	0
	B	0	61	0	46	N/A	N/A
Manitoba	A(H3N2)	0	21	0	16	13	0
	A(H1N1)	0	1	0	1	1	0
	B	0	1	0	1	N/A	N/A

N/A = Not applicable

Table 2. Influenza Strain Characterization reported by NML since September 1, 2017, 2017–2018

Strain	Number of viruses	
	Canada	Manitoba
A/Hong Kong/4801/2014 (H3N2)-like	52	5
A/Michigan/45/2015 (H1N1)-like	20	1
B/Brisbane/60/2008-like	6	0
B/Phuket/3073/2013-like	57	1

Since September 1, 2017, NML has characterized 273 influenza A and B viruses.

1. 160 influenza A(H3N2) viruses:

- 52 influenza A(H3N2) viruses were antigenically characterized as A/Hong Kong/4801/2014, the influenza A(H3N2) component of the 2017-2018 Northern Hemisphere influenza vaccine. In those 32 viruses, 29 belonged to genetic group 3C.2a and 3 belonged to subclade group 3C.2a1.
- 138 influenza A(H3N2) viruses did not grow to sufficient hemagglutination titers for antigenic characterization by hemagglutination inhibition assays. Therefore, genetic characterization was performed. Sequence analysis of the HA gene of these viruses showed that 111 H3N2 viruses belonged to genetic group 3C.2a and 27 belonged to subclade 3C.2a1. A/Hong Kong/4801/2014(H3N2)-like virus, the vaccine strain, belongs to genetic group 3C.2a.

2. 20 influenza A(H1N1) viruses:

- 11 influenza A(H1N1) viruses characterized were antigenically similar to A/Michigan/45/2015, the influenza A(H1N1) component in the vaccine.

3. 63 influenza B viruses:

- 6 influenza B viruses were antigenically similar to B/Brisbane/60/2008-like (Victoria lineage), the influenza B component in the vaccine.
- 57 influenza B viruses characterized were characterized as B/Phuket/3073/2013 (Yamagata lineage), the influenza B component only in the quadrivalent vaccine.

Appendix

Data Sources

Laboratory Surveillance

Detections of influenza nucleic acid detection, culture isolation, and enzyme immunoassay (EIA) are reported from Cadham Provincial Laboratory (CPL) and occasionally other laboratories. These reports are forwarded to Epidemiology and Surveillance (E&S) within 24 hours of confirmation. CPL also performs testing for other respiratory viruses including parainfluenza, RSV, adenovirus, rhinovirus, coronavirus, enterovirus, and bocavirus, which are reported to E&S on a weekly basis.

Influenza Associated Hospitalizations, ICU Admissions and Deaths

Each influenza season on a weekly basis, the Public Health Office in each Regional Health Authority (RHA) is requested to submit a line list of hospitalizations, Intensive Care Unit (ICU) admissions and deaths for laboratory-confirmed influenza patients that were admitted in hospitals in the reporting RHA or deceased as the registered residents of the reporting RHA.

The reason for the hospitalizations, ICU admissions or the cause of deaths does not have to be attributable to influenza. Instead, a temporal association with a positive influenza laboratory result is sufficient due to the requirement for timely reporting. Submissions are cleaned by E&S to remove duplicate submissions for the same patient within the same illness episode. In this report, only Manitoba residents are included.

Outbreak

Outbreaks are reported to E&S by a phone call or email from public health staff within RHAs or from CPL advising the assignment of an outbreak code. CPL submits both positive and negative laboratory results related to outbreaks to E&S. Outbreak investigations are reported from RHAs to E&S by completing an outbreak summary report form electronically through the Canadian Network for Public Health Intelligence (CNPHI) or on paper.

Health Links – Info Santé

Health Links–Info Santé is a 24-hour, 7-days a week telephone information service. It is staffed by registered nurses with the knowledge to provide answers to health care questions and guidance to appropriate care over the phone. When a caller phones Health Links–Info Santé and selects Influenza Service, they are given an option to select information on (1) the groups of individuals who are at an increased risk of serious illness, (2) how to arrange an influenza vaccine, (3) the annual influenza immunization campaign, or (4) the management of influenza and its potential complications.

ILI Visits to Sentinel Physicians

Manitoba participates in *FluWatch*, the Canada's national surveillance system co-ordinated by Public Health Agency of Canada (PHAC), which monitors the spread of influenza and ILI on a year-round basis. *FluWatch* consists of a network of laboratories, hospitals, doctor's offices and provincial and territorial ministries of health. In 2017–2018, there are 19 sentinel physicians recruited throughout Manitoba reporting to *FluWatch* weekly. E&S receives weekly reports from *FluWatch* which present the ILI rate for Manitoba and for each participating sentinel physician. The reporting sentinel physicians are different by week and their reports may not be representative of ILI activity across the province.

Antiviral Dispensing

The units of antiviral drugs, Oseltamivir and Zanamivir, dispensed from community retail pharmacies to Manitoba residents are reported to E&S from Drug Programs Information Network (DPIN) on a weekly basis. Antiviral drugs dispensed to in-patients or through nursing stations could not be included in this report due to lack of data.

Respiratory Visits to Emergency Department

Daily statistics of visits to Emergency Department (ED) that are related to respiratory illness in Winnipeg Regional Health Authority (WRHA) are submitted to E&S weekly. ILL cases are defined as patients whose triage chief complaints contain either of these symptoms: weakness, shortness of breath, cough, headache, fever, cardiac/respiratory arrest, sore throat, and upper respiratory tract infection complaints.

Immunization

Immunization data were extracted from Panorama, the provincial immunization registry. The report includes clients who have received one or more valid doses of the influenza vaccine and who have active registration with Manitoba Health, Seniors and Active Living (MHSAL) at the time that the report is generated (i.e. it is not based on the number of doses in the registry divided by the population of Manitoba). This report coverage does not include the following:

- clients who have moved or passed away and are now inactive in the client registry;
- clients who receive services from public health in Manitoba, but have been entered in the Immunization Registry as inactive clients;
- clients who are not in the registry, such as immigrants and refugees, visiting students, and visitors who are not registered for health services in Manitoba;
- doses that have not been reported to MHSAL.

Antiviral Resistance

Influenza and Respiratory Viruses Section of National Microbiology Laboratory (NML) undertakes enhanced surveillance, investigations, and research on influenza and other respiratory pathogens. A sample of positive influenza specimens isolated by culture is referred from each provincial laboratory to NML for strain characterization and antiviral resistance testing. The aggregate level information is then shared with provinces and territories on a weekly basis.

Circulating Strain

NML antigenically characterizes influenza viruses received from Canadian laboratories year-round. In Manitoba, representative samples of influenza viruses isolated by culture, from different regions, time periods and outbreaks, is referred from CPL to NML for strain characterization.

Immunization Program

As per World Health Organization (WHO), all seasonal quadrivalent influenza vaccines for 2017–2018 in the northern hemisphere contain:

- A/Hong Kong/4801/2014 (H3N2)-like virus (in trivalent vaccine)
- A/Michigan/45/2015 (H1N1)pdm09-like virus (in trivalent vaccine)
- B/Brisbane/60/2008-like virus (in trivalent vaccine)
- B/Phuket/3073/2013-like virus (in quadrivalent vaccine)

For the 2017–2018 influenza season, MHSAL has been allotted the quadrivalent inactivated vaccines (QIV), Fluzone® Quadrivalent (Sanofi Pasteur) and FluLaval Tetra® (GlaxoSmithKline), and the

quadrivalent live attenuated influenza vaccine (QLAIV) FluMist® Quadrivalent (AstraZeneca), as part of the province's Publicly-Funded Seasonal Influenza Immunization Program.

This year, a new seasonal influenza vaccine, Fluzone® High-Dose, is offered to people 65 years of age or older that are living in long-term care facilities (LTCF) in Manitoba. This vaccine is a trivalent inactivated vaccine (TIV) and contains four times the amount of influenza virus antigen per strain compared to the standard-dose influenza vaccine. Therefore, this vaccine is expected to provide better protection against seasonal influenza than the standard-dose vaccine.

Abbreviations

CPL	Cadham Provincial Laboratory
CNPHI	Canadian Network for Public Health Intelligence
E&S	Epidemiology and Surveillance
ED	Emergency Department
ICU	Intensive Care Unit
ILI	Influenza-Like-Illness
LTCF	Long Term Care Facility
MHSAL	Manitoba Health, Seniors and Active Living
NML	National Microbiology Laboratory
PHAC	Public Health Agency of Canada
RHA	Regional Health Authority
RSV	Respiratory Syncytial Virus
WRHA	Winnipeg Regional Health Authority

Explanatory Notes and Definitions

Cumulative Data

Cumulative data include updates to previous weeks; due to reporting delays or amendments, the sum of weekly report totals may not add up to cumulative totals.

Data Extraction Date

Manitoba-specific information contained within this report is based on data confirmed at 11:00 am on the date of data extraction.

Epidemiology Week

Time trends in this report are presented by [epidemiology week](#), a schedule used by the national FluWatch program coordinated by PHAC.

Incidence Rate

Incidence rate measures the frequency that influenza occurs in a population. It is calculated as the total number of new cases this influenza season multiplied by 100,000 and divided by the total population in each region. Regional populations are based on the Manitoba Health Population Report 2016.

ILI in General Population

Acute onset of respiratory illness with fever and cough and with one or more of the following – sore throat, arthralgia, myalgia, or prostration, which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

ILI Outbreak

Schools: Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI.

Hospitals and residential institutions: Two or more cases of ILI within a seven-day period.

Other settings: Two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. workplace, closed communities.

Specimen Collection Date

The date the laboratory specimen was taken is used to assign cases to the epidemiology week in this report. Occasionally, if the specimen collection date is not available, the laboratory report date will be used.

Epidemiology and Surveillance
Manitoba Health, Seniors and Active Living

flusurveillance@gov.mb.ca

(204)786-7335

Other Epidemiology and Surveillance reports

<http://www.gov.mb.ca/health/publichealth/surveillance/index.html>

National Influenza Surveillance

<http://www.phac-aspc.gc.ca/fluwatch/index-eng.php>