

GEOLOGICAL SETTING - AGASSIZ METALLOTECT

The Agassiz MetalloTECT is located in the northern belt of the Lynn Lake greenstone belt and is characterized by a unique and persistent geophysical signature (Fedikow, 1986a) and a distinctive lithological association that consists of high MgO-Ni-Cr basic rocks (termed "picritic basalts"), iron formation, clastic sedimentary rocks (Fedikow and Gale, 1980) and felsic volcanic rocks (Parberry, 1988). The metalloTECT has a strike length of 70 km and extends from the Spider Lake area (Fedikow, 1986b) to the Sheila Lake-Margaret Lake area (Ferreira, 1988). The Nisku, MacLellan and Rianbow Au - Ag, Pb, Zn deposits, the Dot Lake Au deposit and the Farley Lake Au deposit occur along the metalloTECT (Fedikow et al., 1990).

GENERAL STRATIGRAPHY AND LITHOLOGY

Picritic (high MgO-Ni-Cr) basaltic rocks are the most conspicuous lithology of the Agassiz MetalloTECT. Ranges for MgO, Ni and Cr are: 10.21-18.46%, 383-1179 ppm, and 339-2032 ppm, respectively. Picritic basaltic rocks occur throughout the metalloTECT, but decrease in abundance eastward from the MacLellan deposit. Picritic rocks overlie, and are intercalated with, oxide, sulphide and silicate facies iron formation, are intercalated with dark green basaltic rocks, and are in turn overlain by felsic volcanic rocks. Picritic rocks weather a distinctive blue green to dark green and consist of 0.5 to 4 m thick heterolithic, monolithic and flow breccia, pillow breccia, tuff and pillowed flows (Parberry and Fedikow, 1987). Locally, up to 30%, disseminated, subhedral to euhedral magnetite occurs in the picritic volcanic rocks. The rocks of the Agassiz MetalloTECT are bound to the south and north by aluminous basaltic fragmental rocks (Fedikow, 1986b) and minor felsic and mafic intrusions.

Heterolithic breccia contains two to five clast types that in total make up 40 to 80% of the rock. Clasts are commonly subrounded to subangular, amygdaloidal, and may be aphyric or feldspar- and/or amphibole-phyritic. The clasts range in size from 0.5 by 1 cm to 15 by 30 cm. Breccia groundmass is very fine grained and consists almost entirely of chlorite and amphibole with accessory magnetite. Outcrops of fragmental picritic rock that have been strongly folded may contain up to 10%, 1 to 4 mm amphibole porphyroblasts in the matrix and clasts.

Monolithic breccia contains 20 to 50% light green clasts in a very fine grained, dark green, chlorite-amphibole matrix. Clasts are subangular to subrounded, aphanitic to very fine grained, and contain up to 10%, <2 mm plagioclase + quartz amygdaloids. Clasts are generally elongated parallel to foliation and may be a few to several centimetres in length.

Dark green basaltic rocks with distinctive higher contents of MgO, Ni and Cr (5 to 10% MgO and several hundred ppm Ni and Cr) than the aluminous basaltic rocks (average of three samples = 4.4% MgO, 27 ppm Ni and 91 ppm Cr; Syme, 1988) are intercalated with the picritic basaltic rocks. These mafic volcanic rocks occur along the length of the Agassiz MetalloTECT.

Aphyric and quartz-phyric felsic volcanic rocks occur in several locations along the Agassiz MetalloTECT and are commonly associated with picritic rocks. They are most common in the Barrington Lake area (Fedikow et al., 1990). Quartz-phyric felsic rocks have been noted in drill core at the MacLellan deposit and are considered to overlie the picritic rocks (Fedikow, 1986b).

Exposures of banded iron formation (BIF) along the Agassiz MetalloTECT are sporadic. Geophysical data suggest that iron formation is present along most of the metalloTECT. Most BIF observed in the field is oxide facies iron formation, either as chert/quartz/magnetite or chert/quartz-hematite. These units are generally 0.1 to 1.0 m thick, have limited extent, and are interbedded with basaltic volcanic and/or sedimentary rocks. Subpillowed magnetite/chert BIF at Farley Lake contains gold (Briggs and Taylor, 1987). Silicate facies iron formation has been observed in drill core from the MacLellan deposit and contains 5 to 20%, 5 to 10 mm pink garnets in a fine grained, green, chloritic matrix with minor magnetite and lesser amounts of calcite and amphibole. In drill core, thin 1 to 10 mm cherty layers are commonly intercalated with chlorite-rich layers. The silicate facies BIF does not appear to contain sulfides or gold. Sulfide facies iron formation occurs within picritic basalt and clastic sedimentary rocks at the MacLellan deposit. The facies of iron formation consists of 2 to 15 cm thick laminated, gold-bearing disseminated to solid pyrrhotite and pyrite layers that are rhythmically intercalated with botulic and quartz-rich layers. Sphalerite, quartz and calcite occur as accessory minerals. Gagnon (1991) considers these iron sulphide-quartz layers to be deformed quartz veins.

Clastic sedimentary rocks are intercalated with picritic and nonpicritic volcanic rocks and have been referred to as siltstone, calcareous greywacke, and siliceous tuff (Fedikow, 1986b). Exposures are 0.01 to 2.0 m wide. The sedimentary rocks are fine grained, weather white to brown grey, and may contain up to 2%, 1 to 2 mm disseminated subhedral to euhedral magnetite crystals in a quartz-feldspar + siltite groundmass.

Laminated to bedded, reverse and normally graded siltstone has been identified at the MacLellan deposit. The siliceous and/or botulic-rich layers that host the sulfide mineralization and gold may represent either sedimentary rocks or zones of intense alteration arranged concentrically about a shear zone(s) (Fedikow, 1986b).

Other lithologies in the metalloTECT include mafic to intermediate volcanic rocks and small tonalitic, dioritic and gabbroic intrusions.

STRUCTURAL COMPONENT

Rocks in the northern belt of the Lynn Lake greenstone belt have undergone moderate physical deformation. Gilbert et al. (1980) describe the northern belt as consisting of a homoclinal, north-facing sequence of supracrustal rocks; however, Parberry (1988) notes that within the high-Mg (picritic) volcanic rocks, tops are commonly to the south as indicated by pillow tops, pillow breccia and graded bedding. Isoclinal folds probably resulted in both north and south facing top directions. Strike direction are dominantly eastward and dips are steep. Foliations trend mainly east-northeast. A persistent crenulation cleavage (at 244-278°W), which can be measured over a distance of 5 km, occurs within the picritic basalts at the eastern end of the metalloTECT. Picritic rocks that outcrop in the MacLellan deposit area are characterized by the development of mylonitic textures, shear bands, and pseudo-tachylite.

BARRINGTON LAKE-SPIDER LAKE AREA (WEST)

Picritic Basalts

Picritic tuff outcrops sporadically in the Barrington Lake - Spider Lake area and is intercalated with basaltic rock.

Felsic Volcanic Rocks

In the Barrington Lake area felsic volcanic rocks occur along a 200 to 300 m wide zone that has a strike length of several kilometres. The felsic volcanic rocks consist of quartz-phyric tuff and a slightly coarser grained, botulic tuffaceous rock, both of which are intercalated with picritic and nonpicritic volcanic rocks.

Quartz-phyric tuff contains 10 to 30%, 1 to 6 mm diameter, blue to blue grey, augen-shaped, quartz megacrysts. Minor 10 to 20 cm thick, fine grained aphyric units are found within the quartz-ey felsic volcanic rocks. The aphyric units are considered to be felsic dykes.

The botulic tuffaceous rocks are very fine grained and weather buff white to grey brown. They are composed of a quartz-feldspar groundmass with minor botulic and may contain up to 2%, 1 to 2 mm disseminated subhedral to euhedral magnetite. Aducluar amphibole porphyroblasts are locally present and constitute up to 5% of the rock. Discontinuous amphibole-rich layers, 0.5 to 2.0 cm thick, contain 5 to 20% acicular to blocky amphiboles and 0.3 to 3.0 mm pink-orange garnets. Anneldal plagioclase crystals occur in some outcrops. These units are 0.01 to 2.0 m thick.

Clastic Sedimentary Rocks

In the Barrington Lake area rare, thin graded beds of siltstone are intercalated with mafic volcanic rocks. A calc-silicate unit is present east of the Barrington River.

Other

Massive and strongly foliated mafic volcanic rocks that are similar in appearance to picritic basaltic rocks are exposed in a 300 to 500 m wide by 8 km long area. Rock types include heterolithic breccia, flow breccia, tuff, amphibole-phyric basalt and a distinctive two-clast breccia. These rocks have higher than average Cr contents (average of 32 samples = 254 ppm Cr) than the surrounding aluminous basaltic rocks.

Picritic rocks with high Cr and Ni values (up to 1800 ppm and 240 ppm, respectively) occur in drill core from holes drilled in the MacBride Lake area, 5 km ENE of Spider Lake.

ACKNOWLEDGEMENTS:

This project was initiated by M.A.F. Fedikow of Manitoba Energy and Mines. Geological mapping was conducted during the 1987 and 1988 field seasons. Outcrop base maps were supplied by Manitoba Mineral Resources, Ltd. and by Sherritt Gordon Mines, Ltd. Their assistance is greatly appreciated. Field visits by, and discussions with, M.A.F. Fedikow have aided in mapping and in the understanding of the regional aspects of the Agassiz MetalloTECT. B. Barne, G. Foote, T. Schwartz and N. Simpson are thanked for their assistance in the field.

REFERENCES:

Briggs, D.N. and Taylor, M.R.  
 1987: The Farley Lake gold discovery, Lynn Lake, Manitoba. Geological Association of Canada, Program with Abstracts, v. 12, p. 26.  
 Fedikow, M.A.F.  
 1986a: The Agassiz metalloTECT - Spider Lake area. In Manitoba Energy and Mines, Minerals Division, Report of Field Activities, 1986, p. 23-25.  
 1986b: Geology of the Agassiz stratabound Au-Ag deposit, Lynn Lake, Manitoba, Manitoba Energy and Mines, Open File Report, OF85-5, 80 p.  
 Fedikow, M.A.F. and Gale, G.H.  
 1982: Mineral deposit studies in the Lynn Lake area. Agassiz MetalloTECT. In Manitoba Energy and Mines, Minerals Division, Report of Field Activities, 1982, p. 29-31.  
 Fedikow, M.A.F., Parberry, D. and Ferreira, K.J.  
 1990: Agassiz metalloTECT - a regional metallogenic concept. Lynn Lake area, Manitoba: Manitoba Energy and Mines, Minerals Division, Mineral Deposit Thematic Map Series, MAP 99-1.  
 Ferreira, K.J.  
 1986: Geological investigations in the Sheila Lake-Margaret Lake area. In Manitoba Energy and Mines, Minerals Division, Report of Field Activities, 1986, p. 8-12.  
 Gagnon, J.E.  
 1991: Geology, geochemistry and genesis of the Proterozoic MacLellan Au-Ag deposit, Lynn Lake greenstone belt, Manitoba: University of Windsor, M.Sc. thesis (unpublished), 275p.  
 Gilbert, H.P., Syme, E.C. and Zwanig, H.V.  
 1980: Geology of the metavolcanic and volcanoclastic rocks in the Lynn Lake area, Manitoba Energy and Mines, Geological Paper GP80-1, 118 p.  
 Parberry, D.  
 1988: Investigation of volcanic stratigraphy and iron formation occurrences, Lynn Lake area. In Manitoba Energy and Mines, Minerals Division, Report of Field Activities, 1988, p. 12-15.  
 Parberry, D. and Fedikow, M.A.F.  
 1987: Investigations of Agassiz MetalloTECT stratigraphy. In Manitoba Energy and Mines, Minerals Division, Report of Field Activities, 1987, p. 12-16.  
 Syme, E.  
 1988: Geochemistry of metavolcanic rocks in the Lynn Lake Belt, Manitoba Energy and Mines, Geological Services, Geological Report GR84-1, 84p.

ECONOMIC GEOLOGY REPORT ER91-2

Barrington Lake - Spider Lake Area Map #7

