Manitoba
Energy and Mines

ECONOMIC GEOLOGY REPORT ER91-2

Eagle Lake Area – East Map #2

GEOLOGICAL SETTING - AGASSIZ METALLOTECT

The Agassiz Metallotect is located in the northern belt of the Lynn Lake greenstone belt and is characterized by a unique and persistent geophysical signature (Fedikow, 1986a) and a distinctive lithological association that consists of high MgO-Ni-Cr basaltic rocks (termed "picritic basalts"), iron formation, clastic sedimentary rocks, (Fedikow and Gale, 1982) and felsic volcanic rocks (Parbery, 1988). The metallotect has a strike length of 70 km and extends from the Spider Lake area (Fedikow, 1986b) to the Sheila Lake-Margaret Lake area (Ferreira, 1986). The Nisku, MacLellan and Rainbow Au ± Ag, Pb, Zn deposits, the Dot Lake Au deposit and the Farley Lake Au deposit occur along the metallotect (Fedikow *et al*; 1990).

GENERAL STRATIGRAPHY AND LITHOLOGY

Picritic (high MgO-Ni-Cr) basaltic rocks are the most conspicuous lithology of the Agassiz Metallotect. Ranges for MgO, Ni and Cr are: 10.21-18.46%, 393-1 179 ppm, and 939-2 032 ppm, respectively. Picritic basaltic rocks occur throughout the metallotect, but decrease in abundance eastward from the MacLellan deposit. Picritic rocks overlie, and are intercalated with, oxide, sulphide and silicate facies iron formation, are intercalated with dark green basaltic rocks, and are in turn overlain by felsic volcanic rocks. Picritic rocks weather a distinctive blue green to dark green and consist of 0.5 to 4 m thick heterolithic, monolithic and flow breccia, pillow breccia, tuff and pillowed flows (Parbery and Fedikow, 1987). Locally, up to 30%, disseminated, subhedral to euhedral magnetite occurs in the picritic volcanic rocks. The rocks of the Agassiz Metallotect are bound to the south and north by aluminous basaltic fragmental rocks (Fedikow, 1986b) and minor felsic and mafic intrusions.

Heterolithic breccia contains two to five clast types that in total make up 40 to 80% of the rock. Clasts are commonly subrounded to subangular, amygdaloidal, and may be aphyric or feldspar- and/or amphibole-phyric. The clasts range in size from 0.5 by 1 cm to 15 by 30 cm. Breccia groundmass is very fine grained and consists almost entirely of chlorite and amphibole with accessory magnetite. Outcrops of fragmental picritic rock that have been strongly foliated may contain up to 10%, 1 to 4 mm amphibole porphyroblasts in the matrix and clasts.

Monolithic breccia contains 20 to 50% light green clasts in a very fine grained, dark green, chlorite-amphibole matrix. Clasts are subangular to subrounded, aphanitic to very fine grained, and contain up to 10%, <2 mm plagioclase ± quartz amygdules. Clasts are generally elongated parallel to foliation and may be a few to several centimetres in length.

Dark green basaltic rocks with distinctive higher contents of MgO, Ni and Cr (5 to 10% MgO and several hundred ppm Ni and Cr) than the aluminous basaltic rocks (average of three samples = 4.4% MgO, 57 ppm Ni and 91 ppm Cr; Syme, 1985) are intercalated with the picritic basaltic rocks. These mafic volcanic rocks occur along the length of the Agassiz Metallotect.

Aphyric and quartz-phyric felsic volcanic rocks occur in several locations along the Agassiz Metallotect and are commonly associated with picritic rocks. They are most common in the Barrington Lake area (Fedikow *et al*; 1990). Quartz-phyric felsic rocks have been noted in drill core at the MacLellan deposit and are considered to overlie the picritic rocks (Fedikow, 1986b).

Exposures of banded iron formation (BIF) along the Agassiz Metallotect are sporadic. Geophysical data suggest that iron formation is present along most of the metallotect. Most BIF observed in the field is oxide facies iron formation, either as chert/quartz magnetite or chert/quartz-hematite. These units are generally 0.1 to 1.0 m thick, have limited extent, and are interlayered with basaltic volcanic and/or sedimentary rocks. Sulphidized magnetite/chert BIF at Farley Lake contains gold (Briggs and Taylor, 1987). Silicate facies iron formation has been observed in drill core from the MacLellan deposit and contains 5 to 20%, 5 to 10 mm pink garnets in a fine grained, green, chloritic matrix with minor magnetite and lesser amounts of calcite and amphibole. In drill core, thin 1 to 10 mm cherty layers are commonly intercalated with chlorite-rich layers. The silicate facies BIF does not appear to contain sulfides or gold. Sulfide facies iron formation occurs within picritic basalt and clastic sedimentary rocks at the MacLellan deposit. This facies of iron formation consists of 2 to 15 cm thick laminated, gold-bearing disseminated to solid pyrrhotite and pyrite layers that are rhythmically intercalated with biotite- and quartz-rich layers. Sphalerite, quartz and calcite occur as accessory minerals. Gagnon (1991) considers these iron sulphide-quartz layers to be deformed quartz veins.

Clastic sedimentary rocks are intercalated with picritic and nonpicritic volcanic rocks and have been referred to as siltstone, calcareous greywacke, and siliceous tuff (Fedikow, 1986b). Exposures are 0.01 to 2.0 m wide. The sedimentary rocks are fine grained, weather white to brown grey, and may contain up to 2%, 1 to 2 mm disseminated subhedral to euhedral magnetite crystals in a quartz-feldspar ± biotite groundmass.

Laminated to bedded, reverse and normally graded siltstone has been identified at the MacLellan deposit. The siliceous and/or biotite-rich layers that host the sulfide mineralization and gold may represent either sedimentary rocks or zones of intense alteration arranged concentrically about a shear zone(s) (Fedikow, 1986b).

Other lithologies in the metallotect include mafic to intermediate volcanic rocks and small tonalitic, dioritic and gabbroic intrusions.

STRUCTURAL COMPONENT

Rocks in the northern belt of the Lynn Lake greenstone belt have undergone moderate physical deformation. Gilbert *et al*; (1980) describe the northern belt as consisting of a homoclinal, north-facing sequence of supracrustal rocks; however, Parbery (1988) notes that within the high-Mg (picritic) volcanic rocks, tops are commonly to the south as indicated by pillow tops, pillow breccia and graded bedding. Isoclinal folds probably resulted in both north and south facing top directions. Strike directions are dominantly eastward and dips are steep. Foliations trend mainly east-northeast. A persistent crenulation cleavage (at 244°/78°W), which can be measured over a distance of 5 km, occurs within the picritic basalts at the eastern end of the metallotect. Picritic rocks that outcrop in the MacLellan deposit area are characterized by the development of mylonitic textures, shear bands, and pseudotachylite.

EAGLE LAKE AREA - EAST

Picritic Basalts

Picritic rocks are exposed in a 500 m wide by 12 km long discontinuous zone from Eagle Lake to Gordon Lake (Parbery and Fedikow, 1987). Picritic basaltic rocks in the Eagle Lake area consist of monolithic and heterolithic breccia as well as tuff, flow breccia and pillowed flows. Picritic tuff occurs in 3 to 20 cm thick layers.

Felsic Volcanic Rocks

Felsic volcanic rocks in the Eagle Lake area contain 1%, 0.5 to 3 mm anhedral to subhedral quartz phenocrysts and 1%, 1 to 3 mm anhedral garnets in an aphanitic groundmass. Small areas (up to 6 by 15 cm) that contain up to 10% quartz crystals represent either boudinaged layers or elongated clasts. Felsic volcanic rock units are up to 50 m thick and are intruded by 10 to 20 cm thick amphibolitic dykes.

Other

Aluminous heterolithic volcanic breccia and tuff occur to the north of the picritic rocks.

ACKNOWLEDGEMENTS:

This project was initiated by M.A.F. Fedikow of Manitoba Energy and Mines. Geological mapping was conducted during the 1987 and 1988 field seasons. Outcrop base maps were supplied by Manitoba Mineral Resources, Ltd., and by Sherritt Gordon Mines, Ltd. Their assistance is greatly appreciated. Field visits by, and discussions with, M.A.F. Fedikow have aided in mapping and in the understanding of the regional aspects of the Agassiz Metallotect. B. Barrie, G. Foote, T. Schwartz and M. Simpson are thanked for their assistance in the field.

REFERENCES:

Briggs, D.N. and Taylor, M.R.
1987: The Farley Lake gold discovery, Lynn Lake, Manitoba; Geological Association of Canada, Program with Abstracts, v. 12, p. 26.

Fedikow, M.A.F.
1986a: The Agassiz metallotect - Spider Lake area; **in** Manitoba Energy and Mines, Minerals Division, Report of Field Activities, 1986, p. 23-25.
1986b: Geology of the Agassiz stratabound Au-Ag deposit, Lynn Lake, Manitoba; Manitoba Energy and Mines, Open File Report, OF85-5, 80 p.

Fedikow, M.A.F. and Gale, G.H.
1982: Mineral deposit studies in the Lynn Lake area: Agassiz Metallotect; **in** Manitoba Energy and Mines, Minerals Division, Report of Field Activities, 1982, p. 29-31.

Fedikow, M.A.F., Parbery, D. and Ferreira, K.J.
1990: Agassiz metallotect - a regional metallogenic concept, Lynn Lake area, Manitoba; Manitoba Energy and Mines, Minerals Division, Mineral Deposit Thematic Map Series, MAP 89-1.

Ferreira, K.J.
1986: Geological investigations in the Sheila Lake-Margaret Lake area; **in** Manitoba Energy and Mines, Minerals Division, Report of Field Activities, 1986, p. 8-12.

Gagnon, J.E.
1991: Geology, geochemistry and genesis of the Proterozoic MacLellan Au-Ag deposit, Lynn Lake greenstone belt, Manitoba; University of Windsor, M.Sc. thesis (unpublished), 275p.

Gilbert, H.P., Syme, E.C. and Zwanzig, H.V.
1980: Geology of the metavolcanic and volcaniclastic rocks in the Lynn Lake area; Manitoba Energy and Mines, Geological Paper GP80-1, 118 p.

Parbery, D.
1988: Investigation of volcanic stratigraphy and iron formation occurrences, Lynn Lake area; **in** Manitoba Energy and Mines, Minerals Division, Report of Field Activities, 1988, p. 12-15.

Parbery, D. and Fedikow, M.A.F.
1987: Investigations of Agassiz Metallotect stratigraphy; **in** Manitoba Energy and Mines, Minerals Division, Report of Field Activities, 1987, p. 12-16.

Syme, E.
1985: Geochemistry of metavolcanic rocks in the Lynn Lake Belt; Manitoba Energy and Mines, Geological Services, Geological Report GR84-1, 84p.

INDEX MAP



AGASSIZ METALLOTECT PROJECT



SYMBOLS

X	Outcrop		Alteration and Mineralization
△	Frost heaved block	Si	silica
	Iron formation (DDH indicated)	ep	epidote
	DDH(SGM, MMR, Spy, CE, Jit and EL holes)	ch	chlorite
	Portage, trail or drill road	cp	chalcopyrite
	Low/high ground boundary	hm	hematite
	Swamp	kf	K-feldspar
S. P.	Sand plain	mt	magnetite
	Trench	po	pyrrhotite
	Farley Au deposit	py	pyrite
	Geological contact (known, assumed)	fu	fuchsite
70	Bedding (inclined, vertical, dip unknown, tops unknown)	q.v.	quartz vein, inclined, dip unknown
	Pillowed flow (tops known, tops unknown)	r.w.	rusty weathered outcrop rusty weathered zone
70	Foliation (inclined, vertical, dip unknown)	• 03301	sample number (collected in 1987)
75	Fracture set (inclined, vertical)	• 202	sample number (collected in 1988)
60 75 80	Conjugate fractures		
85	Plunge of crenulation fold		

LEGEND

PRE- AND POST-SICKLE GROUP INTRUSIVE ROCKS

11 Fine grained mafic dyke

10 Biotite-quartz felsic dyke

9 Amphibolite

8b Gabbro

8a Diorite

7 Granodiorite

6 Syenite

WASEKWAN GROUP VOLCANIC AND SEDIMENTARY ROCKS

5 Felsic volcanic rocks

5d Felsic dykes
5c Dacite, dacitic tuff breccia, and lapilli tuff
5b Rhyolite breccia
5a Felsic tuff, **5a1** rhyolite tuff and lapilli tuff, **5a2**, quartz-eye tuff, **5a3**, feldspar-quartz crystal tuff

4 Mafic-intermediate volcanic rocks

4f Amphibole and/or plagioclase-phyric basalt, massive mafic to intermediate flows and amygdaloidal mafic flows
4e Mafic to intermediate pillowed flows
4d Mafic to intermediate flow breccia
4c Mafic to intermediate breccia, **4c1** mafic to intermediate heterolithic breccia and heterolithic tuff breccia, **4c2** polymictic conglomerate, **4c3** two-clast breccia

4b Mafic fragmental rocks, **4b1** mafic heterolithic lapilli tuff and mafic lapilli tuff, **4b2** mafic plagioclase crystal tuff and mafic to intermediate crystal tuff, **4b3** mafic to intermediate tuff
4a Intermediate volcanic rock

3 Picritic volcanic rocks

3d Amphibole-phyric pillowed flows, pillowed flows and pillow breccia
3c Flow breccia
3b Breccia, **3b1** heterolithic breccia and heterolithic tuff breccia, **3b2** monolithic breccia, **3b3** two-clast breccia
3a Picritic fragmental rocks, **3a1** heterolithic lapilli tuff, monolithic lapilli tuff, and lapillistone, **3a2** pyroxene-phyric and hornblende-phyric crystal tuff, **3a3** massive tuff and banded tuff

2 Chemical and detrital sedimentary rocks

2e Calc-silicate unit
2d Magnetite-quartz banded iron formation
2c Hematite-quartz banded iron formation
2b Magnetite-chlorite banded iron formation
2a Sedimentary rocks, (miscellaneous) **2a1** laminated siliceous sedimentary rocks, **2a2** greywacke

1 Mafic to intermediate volcanic rocks

1c Amphibole-plagioclase crystal tuff and intermediate amphibole-phyric rocks
1b Mafic to intermediate volcanic rocks and amygdaloidal mafic volcanic rocks
1a Mafic crystal tuff

Scale 1: 5 000

METRES 250 0 250 500 METRES

Geology by D. Parbery

Cartography by C. Wojciechowski, M. Timcoe