

GEOLOGICAL SETTING - AGASSIZ METALLOTECT

The Agassiz MetalloTECT is located in the northern belt of the Lynn Lake greenstone belt and is characterized by a unique and persistent geophysical signature (Fedkow, 1986a) and a distinctive lithological association that consists of high MgO-Ni-Cr basaltic rocks (termed 'picritic basalt'), iron formation, clastic sedimentary rocks (Fedkow and Gale, 1982) and felsic volcanic rocks (Parberry, 1988). The metalloTECT has a strike length of 70 km and extends from the Spider Lake area (Fedkow, 1986b) to the Sheila Lake-Margaret Lake area (Ferreira, 1988). The Nisku, MacLellan and Rainbow Au, Ag, Pb, Zn deposits, the Dof Lake Au deposit and the Farley Lake Au deposit occur along the metalloTECT (Fedkow et al., 1990).

GENERAL STRATIGRAPHY AND LITHOLOGY

Picritic (high MgO-Ni-Cr) basaltic rocks are the most conspicuous lithology of the Agassiz MetalloTECT. Ranges for MgO, Ni and Cr are: 10.21-18.46%, 393.1-179 ppm, and 939.2-932 ppm, respectively. Picritic basaltic rocks occur throughout the metalloTECT, but decrease in abundance eastward from the MacLellan deposit. Picritic rocks overlie, and are intercalated with, oxide, sulphide and silicate facies iron formation, are intercalated with dark green basaltic rocks, and are in turn overlain by felsic volcanic rocks. Picritic rocks weather a distinctive blue green to dark green and consist of 0.5 to 4 m thick heterolithic, monolithic and flow breccia, pillow breccia, tuff and pillowed flow (Parberry et al., 1988). Locally, up to 30% disseminated, subhedral to euhedral magnetite occurs in the picritic volcanic rocks. The rocks of the Agassiz MetalloTECT are bound to the south and north by aluminum basaltic fragmentary rocks (Fedkow, 1986b) and minor felsic and mafic intrusions.

Heterolithic breccia contains two to five clast types that in total make up 40 to 80% of the rock. Clasts are commonly subrounded to subangular, amygdaloidal, and may be phytic or feldspar- and/or amphibole-phyric. The clast range in size from 0.5 by 1 cm to 15 by 30 cm. Breccia groundmass is very fine grained and consists almost entirely of chlorite and amphibole with accessory magnetite. Outcrops of fragmental picritic rock that have been strongly foliated may contain up to 10% of 1 to 4 mm amphibole porphyroblasts in the matrix and clasts.

Monolithic breccia contains 20 to 50% light green clasts in a very fine grained, dark green, chlorite-amphibole matrix. Clasts are subangular to subrounded, aphanitic to very fine grained, and contain up to 10% of 2 mm diopside + quartz amphiboles. Clasts are generally elongated parallel to foliation and may be a few to several centimetres in length.

Dark green basaltic rocks with distinctive higher contents of MgO, Ni and Cr (5 to 10% MgO and several hundred ppm Ni and Cr) than the aluminum basaltic rocks (average of three samples = 4.4% MgO, 57 ppm Ni and 51 ppm Cr; Syme, 1988) are intercalated with the picritic basaltic rocks. These mafic volcanic rocks occur along the length of the Agassiz MetalloTECT.

Andesitic and quartz-phyric felsic volcanic rocks occur in several locations along the Agassiz MetalloTECT and are commonly associated with picritic rocks. They are most common in the Barrington Lake area (Fedkow et al., 1990). Quartz-phyric felsic rocks have been noted in drill core at the MacLellan deposit and are considered to overlie the picritic rocks (Fedkow, 1986b).

Exposures of banded iron formation (BIF) along the Agassiz MetalloTECT are sporadic. Geo-physical data suggest that iron formation is present along most of the metalloTECT. Most BIF observed in the field is oxide facies iron formation, either as chert-quartz magnetite or chert-quartz-hematite. These units are generally 0.1 to 1.0 m thick, have limited extent, and are interlayered with basaltic volcanic and/or sedimentary rocks. Subhedral magnetite-chert BIF at Farley Lake contains gold (Briggs and Taylor, 1987). Silicate facies iron formation has been observed in drill core from the MacLellan deposit and contains 5 to 20%, 5 to 10 mm pink garnets in a fine grained, green, chloritic matrix with minor magnetite and lesser amounts of calcite and amphibole. In drill core, thin 1 to 10 mm cherty layers are commonly intercalated with chlorite-rich layers. The silicate facies BIF does not appear to contain sulphides or goethite. Sulphide facies iron formation occurs within picritic basalt and clastic appeal to contain sulphides or goethite. This facies of iron formation consists of 2 to 15 cm thick laminated, gold-bearing disseminated to subhedral magnetite layers that are rhythmically intercalated with biotite- and quartz-rich layers. Silicified, quartz and calcite occur as accessory minerals. Gagnon (1991) considers these iron sulphide quartz layers to be deformed quartz veins.

Clastic sedimentary rocks are intercalated with picritic and nonpicritic volcanic rocks and have been referred to as siltstone, calcareous greywacke, and siliceous tuff (Fedkow, 1988b). Exposures are 0.1 to 2.0 m wide. The sedimentary rocks are fine grained, weather white to brown grey, and may contain up to 2% of 1 to 2 mm disseminated subhedral to euhedral magnetite crystals in a quartz-hematite-biotite groundmass.

Laminated to bedded, reverse and normally graded siltstone has been identified at the MacLellan deposit. The siliceous and/or biotite-rich layers that host the sulfide mineralization and gold may represent either sedimentary rocks or zones of intense alteration arranged concentrically about a shear zone(s) (Fedkow, 1988b).

Other lithologies in the metalloTECT include mafic to intermediate volcanic rocks and small tonalitic, dioritic and gabbroic intrusions.

STRUCTURAL COMPONENT

Rocks in the northern belt of the Lynn Lake greenstone belt have undergone moderate physical deformation. Gilbert et al. (1988) describe the northern belt as consisting of a horizontal, north-trending sequence of supracrustal rocks; however, Parberry (1988) notes that within the high-Mg picritic volcanic rocks, top to the south as indicated by pillow top, pillow breccia and graded bedding. Localized folds probably resulted in both north and south being top directions. Strike directions are dominantly eastward and dips are steep. Foliations trend mainly east-northeast. A persistent crenulation cleavage (at 244°/78°N), which can be measured over a distance of 5 km, occurs within the picritic basalts at the eastern end of the metalloTECT. Picritic rocks that outcrop in the MacLellan deposit area are characterized by the development of mylonitic textures, shear bands, and pseudotachylite.

BARRINGTON LAKE-SPIDER LAKE AREA (EAST)

Picritic Basalts

Picritic tuff outcrops sporadically in the Barrington Lake - Spider Lake area and is intercalated with basaltic rock.

Felsic Volcanic Rocks

In the Barrington Lake area felsic volcanic rocks occur along a 200 to 300 m wide zone that has a strike length of several kilometres. The felsic volcanic rocks consist of quartz-phyric tuff and a slightly coarser grained, biotite-sulphateous rock, both of which are intercalated with picritic and nonpicritic volcanic rocks.

Quartz-phyric tuff contains 10 to 30%, 1 to 5 mm diameter, blue to blue grey, augen-shaped, felsic volcanic rocks. The aphyric units are considered to be felsic dykes.

The biotite-sulphateous rocks are very fine grained and weather buff white to grey brown. They are composed of a quartz-feldspar groundmass with minor biotite and may contain up to 2%, 1 to 2 mm disseminated subhedral to euhedral magnetite. Acicular amphibole porphyroblasts are locally present and constitute up to 5% of the rock. Discontinuous amphibole-rich layers, 0.5 to 2.0 cm thick, contain 5 to 20% acicular to blocky amphibole and 0.5 to 3.0 mm pink-orange garnets. Amphibolite-plagioclase crystals occur in some outcrops. These units are 0.1 to 2.0 m thick.

Clastic Sedimentary Rocks

In the Barrington Lake area, thin graded beds of siltstone are intercalated with mafic volcanic rocks. A calc-silicate unit is present east of the Barrington River.

Other

Massive and strongly foliated mafic volcanic rocks that are similar in appearance to picritic basaltic rocks are exposed in a 300 to 500 m wide by 8 km long area. Rock types include heterolithic breccia, flow breccia, tuff, amphibole-phyric basalt and a distinctive two-clast breccia. These rocks have higher than average Cr contents (average of 32 samples = 224 ppm Cr) than the surrounding aluminum basaltic rocks.

Picritic rocks with high Cr and Ni values (up to 1800 ppm and 240 ppm, respectively) occur in drill core from holes drilled in the MacBride Lake area, 5 km ENE of Spider Lake.

ACKNOWLEDGEMENTS:

This project was initiated by M.A.F. Fedkow of Manitoba Energy and Mines, Geological mapping was conducted during the 1987 and 1988 field seasons. Outcrop base maps were supplied by Manitoba Mineral Resources, Ltd. and by Sherritt Gordon Mines, Ltd. Their assistance is greatly appreciated. Field visits by, and discussions with, M.A.F. Fedkow have aided in mapping and in the understanding of the regional aspects of the Agassiz MetalloTECT. B. Barne, G. Foot, T. Schwartz and M. Simpson are thanked for their assistance in the field.

ECONOMIC GEOLOGY REPORT ER91-2

Barrington Lake - Spider Lake Area Map #8

