



**LEGEND**

This legend is common to GSC maps 2048A–2060A, and MGS geoscientific maps MAP2003-1–MAP2003-12. Coloured legend blocks indicate map units that appear on this map. Not all map symbols shown in the legend necessarily appear on this map.

**QUATERNARY**

**NONGLACIAL DEPOSITS**

**O** Organic deposits: peat, muck; <1–5 m thick; very low relief wetland deposits; accumulated in fen, bog, swamp, and marsh settings.

**E** Eolian sediments: fine sand; 1–5 m thick; dunes; formed by wind prior to stabilization by vegetation, in most cases on subaqueous outwash sand.

**Lm** Shoreline sediments: sand and gravel; 1–2 m thick; beaches; formed by waves at the margins of modern lakes.

**ALLUVIAL SEDIMENTS:** sand and gravel, sand, silt, clay, organic detritus; 1–20 m thick; channel and overbank sediments; deposited by postglacial rivers.

**Ap** Overbank deposits.

**Ac** Channel deposits.

**GLACIOACUSTRINE DEPOSITS**

**Ls** Shoreline deposits.

**Li** Littoral deposits.

**Lz** Clayey to sandy silt.

**Lc** Clay to silty clay.

**GLACIOFLUVIAL DEPOSITS**

**Gs** Subaqueous outwash: fine sand, minor gravel, thin silt and clay interbeds; 1–75 m thick; subaqueous outwash fans; deposited near the ice margin in glacial Lake Agassiz by meltwater turbidly currents, commonly reshaped by wave erosion and reworked by wind.

**ICE-CONTACT GLACIOFLUVIAL SEDIMENTS:** sand and gravel; 1–20 m thick; complex deposits, belts with single or multiple esker ridges and kames, as well as thin, low-relief deposits; deposited in contact with glacial ice by meltwater.

**Gc** Predominantly derived from carbonate rocks.

**Gp** Predominantly derived from igneous and metamorphic rocks.

**GLACIAL DEPOSITS**

**T** Till: calcareous silt diamiction; 1–75 m thick; low-relief, commonly streamlined deposits; subglacial deposits; largely derived from carbonate rocks; thicker sequences consist of multiple units of varying texture, commonly scoured by icebergs; covered discontinuously by thin veneers (<1 m) of glaciolacustrine and glaciofluvial sediments.

**Tc** Predominantly derived from carbonate rocks.

**Tp** Predominantly derived from igneous and metamorphic rocks.

**DISCONTINUOUS TILL AND ASSOCIATED GLACIOFLUVIAL SEDIMENTS:** gravely silt to sand diamiction, sand and gravel; 1–20 m thick; low-relief deposits between bedrock outcrops making up 25–75% of the area; sandy till interbedded and interspersed with nearly equal and often greater amounts of sandy glaciofluvial sediments, as well as minor glaciolacustrine sediments.

**PRE-QUATERNARY**

**ROCK:** >75% bedrock outcrop; Paleozoic carbonate-dominated rocks in areas west and south of Lake Winnipeg, exposed typically as glacially striated, low-relief surfaces; in Precambrian terrane, generally unweathered intrusive, metasedimentary, and metavolcanic rocks having a glacially scoured irregular surface with high local relief; includes patches of thin glacial sediments and organic material.

**Rc** Paleozoic sedimentary rocks.

**Rp** Precambrian igneous and metamorphic rocks.

Geological boundary (approximate) .....  
Built-up area (map GSC 2055A / MGS MAP2003-7 only) .....  
Mine waste .....  
Peat-extraction area .....  
Gravel pit .....  
Mine or bedrock quarry .....  
Stabilized dunes .....  
Abandoned channel .....  
Minor beach ridge .....  
Wave-cut scarp .....  
Groundwater seeping channel .....  
Piping depression .....  
Iceberg scour .....  
Tunnel valley .....  
Esker (direction of flow indicated) .....  
Streamlined landform .....  
Glacial striae .....  
Crossed striae (numbers indicate relative age, 1 being the oldest) .....  
Small bedrock outcrop .....

Geology by G.L.D. Matile, Manitoba Geological Survey, 1993–1994

Co-ordinated by H. Threlkoff and G.L.D. Matile through the auspices of the Southern Plains NATMAP Project and the Winnipeg Region NATMAP Project

Mapping was extended into the State of Minnesota with the permission of the Director, Minnesota Geological Survey

Digital cartography by P.A. Melbourne, Earth Sciences Sector Information Division (ESS Info)

GSC MAP 2060A  
MGS GEOSCIENTIFIC MAP MAP2003-12  
SURFICIAL GEOLOGY  
**WHITEMOUTH LAKE**  
MANITOBA–ONTARIO–MINNESOTA

Scale 1:100 000/Échelle 1/100 000

Universal Transverse Mercator Projection  
North American Datum 1983  
© Her Majesty the Queen in Right of Canada 2004

Projection transversale universelle de Mercator  
Système de référence géodésique nord-américain, 1983  
© Sa Majesté la Reine du chef du Canada 2004

This map was produced from processes that conform to the ESS Info Publishing Services Subdivision Quality Management System, registered to the ISO 9001:2003 standard

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada and the Manitoba Geological Survey

Digital base map from data compiled by Geomatics Canada, modified by ESS Info

Mean magnetic declination 2004, 3°08' E, decreasing 5.6' annually. Readings vary from 0°40' E in the SW corner to 2°36' E in the NE corner of the map.

Elevations in feet above mean sea level

02 413	02 414	02 415	02 416	02 417	02 418
GSC 2048A	GSC 2049A	GSC 2050A	GSC 2051A	GSC 2052A	GSC 2053A
MGS MAP2003-1	MGS MAP2003-2	MGS MAP2003-3	MGS MAP2003-4	MGS MAP2003-5	MGS MAP2003-6
GSC 2054A	GSC 2055A	GSC 2056A	GSC 2057A	GSC 2058A	GSC 2059A
MGS MAP2003-7	MGS MAP2003-8	MGS MAP2003-9	MGS MAP2003-10	MGS MAP2003-11	MGS MAP2003-12
GSC 2060A	GSC 2061A	GSC 2062A	GSC 2063A	GSC 2064A	GSC 2065A
MGS MAP2003-13	MGS MAP2003-14	MGS MAP2003-15	MGS MAP2003-16	MGS MAP2003-17	MGS MAP2003-18

NATIONAL TRANSVERSE MERCATOR SYSTEM REFERENCE AND INDEX TO SURFICIAL GEOLOGICAL SURVEY OF CANADA AND MANITOBA GEOLOGICAL SURVEY MAPS

Recommended citation:  
Matile, G.L.D., 2004. Surficial geology, Whitemouth Lake, Manitoba–Ontario–Minnesota. Geological Survey of Canada, Map 2060A; Manitoba Industry, Economic Development and Mines, Manitoba Geological Survey, Geoscientific Map MAP2003-12, scale 1:100 000.