




Geology of the Hidden and Louis formations in the Louis Lake area, Flin Flon region, Manitoba and Saskatchewan (part of NTS 63K13)



LEGEND


- <1.845 Ga INTRUSIVE ROCKS**
 - B1** Boundary intrusions; fine- to coarse-grained gabbro, locally with various xenoliths
 - Z1** Phantom Lake intrusions
 - G1** Granodiorite sills and dikes (?)
- MISSILI GROUP**
 - X1** Conglomerate, wacke, sandstone and siltstone
- FLIN FLON ARC ASSEMBLAGE (>1.88 Ga ROCKS)**
 - Synvolcanic intrusive rocks**
 - D6** Gabbroic textured, medium- to coarse-grained dikes and sills
 - D5** Gabbroic textured, fine- to medium-grained dikes and sills
 - D4** Aphyric felsic dikes and sills
 - D3a** Fine-grained, grey andesitic dikes with 5 to 20%, 2 to 7 mm amphibole crystals
 - D3b** Coarse-grained, grey, plagioclase-rich andesitic dikes
 - D2a** Plagioclase (5 to 15%, 2 to 10 mm) and pyroxene (1 to 15%, 2 to 10 mm) fine- to medium-grained porphyritic mafic dikes and sills
 - D2b** Pyroxene (5 to 8%, 2 to 10 mm) fine- to medium-grained porphyritic mafic dikes and sills
 - D2c** Plagioclase (5 to 30%, 3 to 15 mm) with 5 to 15% quartz amygdaloid, fine- to medium-grained porphyritic mafic dikes and sills
 - D1** Aphyric to sparsely plagioclase phyric (<5%), fine to medium grained, massive, commonly flow-banded and commonly containing quartz amygdaloid (<5%)
 - Louis formation**
 - Undivided volcanic rocks**
 - L3** Aphyric, pillowed to massive basalt flows
 - L2** Mafic volcanoclastic rocks
 - a) mafic lapillistone
 - b) finely laminated mafic tuff
 - L1** Plagioclase (>15%) and pyroxene (>5%) phyric basalt flows
 - a) massive
 - b) pillowed
 - c) contains included tuff
 - d) in situ breccia
 - Icehouse member**
 - C2** Mafic volcanoclastic rocks
 - a) heterolithic lapillistone, containing plagioclase- and pyroxene-porphyritic, non-amygdaloidal to amygdaloidal basalt clasts (70%), aphyric non-amygdaloidal to amygdaloidal (5-10%) basalt clasts, minor (<5%) aphyric rhyolite clasts in a plagioclase and pyroxene crystal-rich mafic tuff
 - b) finely laminated mafic tuff
 - C1** Strongly plagioclase (>25%) and pyroxene (>15%) porphyritic basalt flows
 - a) massive
 - b) pillowed with tuff between pillows
 - Tower member**
 - T2** Mafic tuff to mafic lapilli-tuff with aphyric rhyolite clasts
 - T1** Massive to in situ brecciated aphyric to sparsely plagioclase (<5%) phyric rhyolite
 - Hidden formation**
 - Undivided intrusive rocks**
 - H1** Pyroxene-phyric basaltic cryptoflows and shallow intrusions
 - a) massive, 1 to 5% pyroxene phenocrysts
 - b) contains included tuff
 - c) peperitic
 - 1920 unit**
 - I1** Amygdaloidal amphibole-phyric andesite cryptoflows
 - a) massive cryptoflows, 1 to 3 mm amphibole phenocrysts
 - b) pillowed cryptoflow, 1 to 3 mm amphibole phenocrysts
 - c) contains included tuff
 - d) peperitic
 - Stockwell member**
 - N2** Mafic heterolithic lapillistone, containing aphyric non-amygdaloidal to amygdaloidal (1-80%) basalt clasts, minor (<1%) felsic clasts
 - N1** Plagioclase (>20%) and pyroxene (<5%) phyric basalt cryptoflows and flows with local peperite
 - a) massive
 - b) pillowed
 - c) contains included tuff
 - d) peperitic
 - Reservoir member**
 - R4** Mafic volcanoclastic rocks
 - R3** Massive plagioclase (>15% to <20%) and pyroxene (>15% to <20%) porphyritic basalt flows
 - R2** Plagioclase (>5% to <10%) phyric basaltic flows
 - a) massive, may contain up to 5% feldspar phenocrysts, quartz amygdaloidal
 - b) pillowed, may contain up to 5% feldspar phenocrysts, quartz amygdaloidal
 - c) in situ breccia
 - R1** Aphyric to weakly (<5%) plagioclase-phyric basaltic cryptoflows and flows with local peperite
 - a) massive, may contain up to 5% feldspar phenocrysts, quartz amygdaloidal
 - b) pillowed, may contain up to 5% feldspar phenocrysts, quartz amygdaloidal
 - c) contains included tuff
 - d) peperitic
 - e) in situ breccia
 - Undivided volcanoclastic rocks**
 - V2** Mafic volcanoclastic rocks
 - V1** Felsic volcanoclastic rocks, pumice-bearing
 - Other**
 - GS** Gossan patches
 - E1** Regolith: weathered profile of aphyric basalt (Hidden formation)

NOTE: units with greyed-out text do not appear on map

SYMBOLS

- Contact: defined, approximate, assumed
- Facies: defined, approximate, assumed
- Fault: defined, approximate, assumed
- Thrust fault: defined, approximate, assumed
- Syncline fold axis
- Mapping limit
- Outcrop
- Bedding: tops unknown, known
- Foliation: generation unknown, 1st, 2nd
- Flow contact: tops known
- Joint
- Vein
- L-fabric: generation unknown, 1st
- Paved road
- Gravel road
- Trail
- Provincial boundary


Index map



Geology by: Y. M. DeWolfe¹
 Digital cartography by: M. E. McFarlane

Published by Manitoba Science, Technology, Energy and Mines, Manitoba Geological Survey, 2007
 This map is available to download free of charge at manitoba.ca/minerals

¹ Mineral Exploration Research Centre, Department of Earth Sciences, Laurentian University, 933 Ramsay Lake Road, Sudbury, Ontario, P3E 6C7



Suggested reference:
 DeWolfe, Y.M., 2007: Geology of the Hidden and Louis formations in the Louis Lake area, Flin Flon region, Manitoba and Saskatchewan (part of NTS 63K13); Manitoba Science, Technology, Energy and Mines, Manitoba Geological Survey, Open File OF2007-4, map at 1:2000 scale.



1:2000

