



# GEOLOGY OF THE ORDOVICIAN / SILURIAN STONEWALL FORMATION IN MANITOBA

## Stratigraphic Map Series Os - 1

### STONEWALL FORMATION

#### Geological Framework

The Stonewall Formation is relatively uniform in thickness and lithology throughout southwestern Manitoba, and reflects a relative stabilization of the tectonic framework in latest Ordovician time. In many ways, the Stonewall Formation seems more closely related to the Silurian stable shelf deposits than to the more basinal Ordovician strata. Although, the above-noted regional northward thinning affects all Ordovician units except the Upper Stonewall.

The sparsely fossiliferous dolomites of the Stonewall Formation, seen at the type section at Stonewall Quarry Park, represent the uppermost beds of the Ordovician succession. The complete Stonewall section, as defined in the subsurface, includes a lower and an upper unit separated by a thin sandy argillaceous marker (i-zone of Porter and Fuller, 1959). Only the lower unit is seen at the Stonewall Quarry. Porter and Fuller (1959) and Brindle (1960) have suggested that the mid-Stonewall marker (m-marker in Manitoba) may, in fact, mark the Ordovician/Silurian boundary and this position has been adopted for this map. This marker, if present in the quarry, would occur only a few feet (half a metre) above the top of the quarry section. The type Williston Member (present in the lower part of the formation) is exposed in a pit at this quarry, and the unit maintains a relatively uniform lithology as an argillaceous dolomite throughout the outcrop belt.

The position of the Ordovician-Silurian boundary in the Williston Basin has been traditionally placed at the i-marker within the upper part of the Stonewall Formation (Porter and Fuller, 1959; Brindle, 1960; McCabe, 1988). Recent biostratigraphic studies confirm this as an approximate placement, both in outcrop (Cormorant Hill roadcut) (Bezys, 1991) and in subsurface (Estehazy 3SWD well, 16-26-20-33WPM, Haidl, 1991; corohole M-1-86, Warren, MB, McCabe, 1986; Norford et al., 1998).

The relatively monotonous dolomite sequence is interrupted only by a number of thin sandy argillaceous marker beds that are believed to represent para-time-stratigraphic markers, minor depositional hiatuses that are very persistent and can be traced throughout most of the Williston Basin area (Porter and Fuller, 1959; King, 1964). The uniformity of Silurian lithology and the persistence of the marker beds attest to the tectonic stability during Silurian time.

Detailed maps have not been compiled for subunits within the Silurian, because of correlation uncertainties. Available data suggest that the tectonic framework was relatively stable throughout Silurian time, with almost no evidence of basin differentiation in the Manitoba portion of the basin. This is evidenced also by the relative lithologic uniformity of the Silurian strata, which consist almost entirely of micritic to intraclastic and stromatolitic dolomites with scattered fossil-fragmental interbeds. These strata are generally representative of deposition under shallow-water, in part slightly restricted conditions, and have been interpreted by Roehl (1967) and others as intertidal to supratidal deposits. In the central part of the basin, the uppermost Silurian beds, stratigraphically above the beds comprising the Manitoba sequence, consist of dolomites with brecciated textures, desiccation cracks, fenestral fabrics, dolomite cements and erosion surfaces (=Upper Interlake). These features indicate periodic subaerial exposure and vadose diagenesis of carbonate deposited under marine or fresh water conditions (Roehl, 1967; Megatham, 1987; Haidl, 1987).

One of the principal post-Paleozoic anomalies relates to the regional configuration of the pre-Jurassic erosion surface. If an east-west structural projection is drawn at the base of the Amaranth Evaporite (the lowest Mesozoic unit that approximates a time-stratigraphic marker), the structure is seen to be rather uniform as far east as the present erosional edge of the unit. However, when the elevations of known Jurassic outcrops and embayments to the east (e.g. Lake St. Martin) are plotted, they are found to fall markedly below the regional projection, and indicate a pronounced structural flexure (perhaps even a structure reversal) at a point east of the main Mesozoic erosional edge. If these outcrops are representative of regional Mesozoic structure, they indicate that the pre-Mesozoic erosion surface approximately paralleled present Paleozoic structure. Surprisingly, no evidence can be seen of any structural flexure in the Paleozoic sequence. Possibly the flexure is masked by the superimposed regional gradient, or possibly this represents a case of structural flexing and later reversal related to the Churchill Superior Boundary Zone.

If the suggested mechanism of differential uplift and subsidence related to the boundary zone is valid, and the pre-Mesozoic unconformity surface developed during a period of maximum uplift, the regional structural profile on the unconformity surface should be "normal" only for the tectonically positive setting. Later reversal or normalization of the tectonic framework would have caused a negative deflection of the erosion surface, which is what seems to be evidenced by the structurally low Mesozoic outcrops.

#### Stratigraphy

The relative uniformity of the Silurian succession, combined with a large area of no outcrop between the northern Grand Rapids area and the southern Fisher Branch area, have resulted in some miscorrelations and resultant problems in stratigraphic nomenclature. The present stratigraphic subdivision of the Silurian outcrop belt is that proposed by Stearn (1956), with slight modification. This detailed subdivision is applicable only to the Manitoba outcrop belt, and cannot satisfactorily be extended to the subsurface. Thus the correlation problems have little stratigraphic significance other than in the outcrop belt itself. The problem arises, however, that the Silurian faunal succession, as derived from the outcrop belt by Stearn, has incorporated some stratigraphic errors so that the faunal elements, in some instances, have been misplaced. A detailed corohole program initiated in 1980 was designed to clarify the correlation problems (McCabe, 1980). These results indicate that the type "Inwood Formation" of the southern area is stratigraphically higher than reported by Stearn, and in fact, may be correlative with the Moose Lake/Atikameg formations of the northern area. Other so-called Inwood and Fisher Branch outcrops in the southern Interlake may also not be stratigraphically consistent. Due to these correlation problems, the term "Inwood" has been dropped (Lammers, 1988).

Stearn (1956) also noted a number of faunal anomalies or inconsistencies in his paleontological studies, and preliminary indications are that most of these inconsistencies can be resolved by revision of the stratigraphic correlations.

In Manitoba, the Stonewall Formation is quite uniform in thickness. It ranges in thickness between 10 to 30 m, with the maximum thickness in southwestern Manitoba.

#### Economics

One oil and gas show has been reported in 3-9-4-11W (Sweet Grass Pilot Mound #1). Crushed stone from the Stonewall Formation has been used as a source of lime from the Stonewall Quarry (now abandoned).

#### References and Selected Bibliography

Andrichuk, J.M. (1959) Ordovician and Silurian stratigraphy and sedimentation in southern Manitoba; American Association of Petroleum Geologists Bulletin, v. 42, No. 10, p. 2333-2398.  
Baillie, A.D. (1951) Silurian geology of the Interlake area, Manitoba; Manitoba Mines Branch, Publication 50-1, 82p.  
Baillie, A.D. (1952) Ordovician geology of Lake Winnipeg and adjacent areas, Manitoba; Manitoba Mines Branch, Publication 51-6, 64p.  
Bannatyne, B.B. (1988) Dolomite resources of southern Manitoba; Manitoba Energy and Mines, Geological Services, Economic Geology Report ER85-1, 39p.  
Bezys, R.K. (1991) Stratigraphic mapping (NTS 63F, 63K) and corohole program 1991; Manitoba Energy and Mines, Minerals Division, Report of Activities 1991, p. 61-73.  
Bezys, R.K. and McCabe, H.R. (1996) Lower to Middle Paleozoic Stratigraphy of Southwestern Manitoba - Field Trip Guidebook B4; Geological Association of Canada/Mineralogical Association of Canada Annual Meeting, Winnipeg, MB, May 27-29, 1996, 92p.  
Brindle, J.E. (1960) The faunas of the Lower Paleozoic carbonate rocks in the subsurface of Saskatchewan; Saskatchewan Department of Mineral Resources, Report No. 52, 45p.  
Cowan, J. (1971) Ordovician and Silurian stratigraphy of the Interlake area, Manitoba; Geological Association of Canada, Special Paper No. 9, p. 235-241.

Haidl, F.M. (1991) Note on the Ordovician/Silurian boundary in southeastern Saskatchewan; Saskatchewan Energy and Mines, Miscellaneous Report 91-4, p. 205-207.  
Johnson, M.E. and Leszczynski, H.L. (1986) Depositional dynamics of cyclic carbonates from the Interlake Group (Lower Silurian) of the Williston Basin; Palaios, v. 1, p. 111-121.  
Kindle, E.M. (1914) The Silurian and Devonian section of western Manitoba; Geological Survey of Canada, Annual Summary Report 1912, p. 247-261.

Kreis, L.K. and Haidl, F.M. (1994) Computer-generated Regional Geological Maps of Southern Saskatchewan: Stonewall Formation; In Summary of Investigations 1994, Saskatchewan Geological Survey, Saskatchewan Energy and Mines, Miscellaneous Report 94-4.  
Lammers, G.E. (1988) Silurian stratigraphy of the Interlake area; Manitoba Energy and Mines, Minerals Division, Report of Field Activities 1988, p. 139-142.

Manitoba Energy and Mines (1979) Geological Map of Manitoba; Map 79-2, 1:1 000 000.  
Manitoba Energy and Mines (1990) Bedrock Geology Compilation Map Series, Preliminary Edition, NTS 63K, 1:250 000.  
Manitoba Energy and Mines (1993) Bedrock Geology Compilation Map Series, Preliminary Edition, Wukusko Lake, NTS 63J, 1:250 000.  
Manitoba Energy and Mines (1994) Geological Highway Map of Manitoba, 1:1 000 000.


Manitoba Energy and Mines (1997) Bedrock Geology Compilation Map Series, Selkirk, NTS 63I, 1:250 000.  
Manitoba Energy and Mines (1998) Lower Paleozoic drill stem tests and oil and gas shows: Three Forks to Precambrian; Petroleum and Energy Division, 29p.  
McCabe, H.R. (1967) Tectonic framework of Paleozoic formations in Manitoba; Transactions of the Canadian Institute of Mining and Metallurgy, v. 70, p. 180-189.

McCabe, H.R. (1980a) Stratigraphic mapping and corohole program, southwest Manitoba; Manitoba Energy and Mines, Report of Field Activities 1980, p. 70-73.  
McCabe, H.R. (1980b) Ordovician - Stonewall Formation (structure contour and isopach maps); Manitoba Stratigraphic Map Series Os-1; Manitoba Energy and Mines, Mineral Resources Division, 1:1 000 000.  
McCabe, H.R. (1988) Preliminary report on Ordovician-Silurian boundary rocks in the Interlake area, Manitoba, Canada; Bulletin of the British Museum of Natural History (Geology), v. 43, p. 255-257.


Norford, B.S., Haidl, F.M., Bezys, R.K., Cecile, M.P., McCabe, H.R., and Paterson, D.F. (1994) Middle Ordovician to Lower Devonian strata of the Western Canada Sedimentary Basin; in Geological Atlas of the Western Canada Sedimentary Basin, G.D. Mossop and I. Shetson (compilers), Canadian Society of Petroleum Geologists and Alberta Research Council, Calgary, p. 109-127.  
Norford, B.S., Nowlan, G.S., Haidl, F.M., and Bezys, R.K. (1998) The Ordovician - Silurian boundary interval in Saskatchewan and Manitoba; in Proceedings of the 8th International Williston Basin Symposium, J.E. Christopher (Ed.), Saskatchewan Geological Society Special Publication.

Porter, J.W. and Fuller, J.G.C.M. (1959) Lower Paleozoic rocks of the northern Williston Basin and adjacent areas; American Association of Petroleum Geologists Bulletin, v. 43, No. 1, p. 124-189.  
Roehl, P.O. (1967) Stony Mountain (Ordovician) and Interlake (Silurian) facies analysis of recent low-energy marine and subaerial carbonates; American Association of Petroleum Geologists Bulletin, v. 51, No. 10, p. 1979-2032.

Smith, D.L. (1963) A lithologic study of the Stony Mountain and Stonewall formations in southern Manitoba; University of Manitoba M.Sc. Thesis, 219p.  
Stearn, C.W. (1956) Stratigraphy and paleontology of the Interlake Group and Stonewall Formation in southern Manitoba; Geological Survey of Canada; Memoir 281, 162p.




Isopach Map  
1 : 2 000 000  
0 25 50 75 100 km




Structure Contour Map  
1 : 2 000 000  
0 25 50 75 100 km

- LEGEND (1 : 2 000 000 maps)
- Stonewall outcrop belt
- Stonewall subcrop belt
- Control well \*
- Isopach Map contour interval (5 m)
- Structure Map contour interval (50 m) (sea-level datum)
- A A' Stratigraphic cross section
- Churchill Superior Boundary Zone



### Location Map



Geology by: R.K. Bezys  
Compilation by: R.K. Bezys and G.G. Conley  
Cartography by: M.E. McFarlane

Suggested reference to this publication:  
Bezys, R.K. and Conley, G.G. 1998: Geology of the Ordovician / Silurian Stonewall Formation in Manitoba; Manitoba Energy and Mines, Stratigraphic Map Series, Os-1, 1:2 000 000.

\* Both confidential and non-confidential wells were used in the construction of these maps; only non-confidential wells are depicted.