

WATER AVAILABILITY AND DROUGHT CONDITIONS REPORT

July 7, 2014

Synopsis/Overview

This Water Availability and Drought Conditions Report provides an update on meteorological and hydrologic drought conditions for Manitoba as of the end of June 2014.

Precipitation indicators show moderately dry conditions prevailed for the Hayes River Basin and area surrounding Norway House.

Monthly stream flow indicators for June indicate flows are normal or above normal for almost all rivers across the province. The Boyne River experienced moderately dry hydrological conditions.

Manitoba Agriculture, Food and Rural Development reported that all dugouts across agri-Manitoba are full or close to full.

Water supply reservoirs in southern and western Manitoba are at or above full supply levels.

Outlook

Environment Canada's seasonal forecast for the next three months (July-August-September, 2014) is for above normal temperatures and below normal precipitation for the entire province (Attachment 4).

Drought Indicators

Two types of drought indicators are assessed across Manitoba - precipitation and stream flow. The indicators describe the severity of dryness in a watershed.

Precipitation is assessed to determine the severity of meteorological dryness and is an indirect measurement of agricultural dryness. Three precipitation indicators are calculated to represent the long term (twelve months), medium term (three months) and short term (one month). Long term and medium term indicators provide the most appropriate assessment of dryness as the short term indicator is influenced by significant rainfall events and spatial variability in rainfall, particularly during summer storms.

The stream flow indicator is used to determine the severity of hydrological dryness in a watershed.

Precipitation Indicators

Precipitation indicators are summarized by basin in Table 1 and on maps in Attachment 1.

Over the long term (twelve months), conditions were normal or above normal throughout the province with the exception of the area between Norway House, Island Lake and Berens River where moderately dry conditions prevailed.

Over the medium term (three months), conditions were normal or above normal throughout the province with the exception of the areas west of Norway House and Gillam where moderately dry conditions prevailed.

Over the short term (one month), conditions were normal or above normal throughout the province with the exception of the areas west of Norway House and Gillam where severely to moderately dry conditions prevailed.

Stream Flow Indicators

Stream flow indicators are summarized by basin in Table 1 and on a map in Attachment 2.

Except for the Boyne River, the monthly stream flow indicators for indicate flows are normal or above normal for all rivers across the province. The Boyne River experienced moderately dry hydrological conditions.

Water Availability

Reservoir Conditions

Water supply reservoirs in southern and western Manitoba are at full or above full supply level (Attachment 3). Provincial water supply reservoirs should have sufficient water supplies for the balance of the year.

On Farm Water Supply

Manitoba Agriculture, Food and Rural Development reports that water levels in dugouts were full, or close to full, in most regions of agri-Manitoba.

Aquifers

Groundwater levels in major aquifers are generally good. Water level responses to seasonal or yearly precipitation fluctuations in most aquifers lag considerably behind surface water responses, so even prolonged periods of below normal precipitation may not have a significant negative effect on groundwater levels. Most aquifers also store very large quantities of groundwater and can continue to provide water during extended periods of dry weather. Consequently, the major concern regarding groundwater and dry periods relates to water levels in shallow wells constructed in near surface sand aquifers. As the water table drops, there is less available drawdown in shallow wells and some wells may 'go dry'.

Forest and Grassland Fires

The Provincial Fire Program reports some large fires in northern Manitoba. More detailed information on fire conditions is available on the Manitoba Conservation and Water Stewardship website under the Fire Program (website <http://www.gov.mb.ca/conservation/fire/>).

Potential Impacts

With Environment Canada's outlook for the next three months for above normal temperature and below normal precipitation, dry conditions may be alleviated. In existing dry areas, particularly the Hayes River basin, there is a risk of grass and forest fire.

Water supply reservoirs are at full or above full supply levels and should have sufficient water supplies for the balance of the year unless prolonged dry conditions occur.

Table 1: Drought Indicators by Major River Basin (Attachments: 1, 2 and 5)

Basin (in Manitoba)	Drought Indicators			
	Monthly Precipitation Indicator (Percent of 1 month Median) June 2014	Monthly Precipitation Indicator (Percent of 3 month Median) (April - June 2014)	Monthly Precipitation Indicator (Percent of 12 month Median) (July 2013- June 2014)	Monthly Flow Indicator June 2014
Red River	Normal conditions	Normal conditions	Normal conditions	Normal conditions except moderately hydrological conditions for the Boyne near Carman
Winnipeg River	Normal conditions	Normal conditions	Normal conditions	Normal conditions
Assiniboine River-Souris River	Normal conditions	Normal conditions	Normal conditions	Normal conditions
Lake Manitoba	Normal conditions	Normal conditions	Normal conditions	Normal conditions
Lake Winnipeg	Normal conditions except moderately dry for the most northeast portion of the basin	Normal conditions	Normal conditions except moderately dry for the northeast portion the basin	Normal conditions
Saskatchewan River	Normal conditions	Normal conditions	Normal conditions	Normal conditions
Nelson River	Normal conditions except moderately to severely dry for the eastern part of the basin	Normal conditions except for moderately dry for eastern parts of the basin	Normal conditions except moderately dry for areas surrounding Norway House	Normal conditions
Hayes River	Moderately to severely dry	Moderately dry conditions	Normal except moderately dry conditions for the southern part of the basin	Normal conditions
Churchill River	Normal conditions	Normal conditions	Normal conditions	Normal conditions
Seal River	Normal conditions	Normal conditions	Normal conditions	Normal conditions

Acknowledgements

This report was prepared with information from the following sources which are gratefully acknowledged:

- Manitoba Infrastructure and Transportation: Flow information:
http://www.gov.mb.ca/mit/floodinfo/floodoutlook/river_conditions.html
- Environment Canada: Flow and Lake information
http://www.wateroffice.ec.gc.ca/index_e.html
- Fire Hazard: <http://www.gov.mb.ca/conservation/fire/>
- Environment Canada 3 month climatic outlook:
http://weatheroffice.gc.ca/saisons/index_e.html
- Manitoba Agriculture, Food and Rural Development:
<http://www.gov.mb.ca/agriculture/crops/seasonal-reports/crop-report-archive/index.html>
- Manitoba Conservation and Water Stewardship Fire Program

For further information, please contact: Mark Lee, Surface Water Management Section, Manitoba Conservation and Water Stewardship, 204-945-5606.

Definition of drought

Meteorological Drought is generally defined by comparing the rainfall in a particular place and at a particular time with the average rainfall for that place. Meteorological drought leads to a depletion of soil moisture and this almost always has an impact on agricultural production. Meteorological droughts only consider the reduction in rainfall amounts and do not take into account the effects of the lack of water on water reservoirs, human needs or on agriculture. A meteorological drought can occur without immediately impacting streamflow, groundwater, or human needs. If a meteorological drought continues, it will eventually begin to affect other water resources.

Agricultural Drought occurs when there is not enough water available for a particular crop to grow at a particular time. Agricultural drought depends not only on the amount of rainfall but also on the use of that water. Agricultural droughts are typically detected after meteorological drought but before a hydrological drought. If agricultural drought continues, plants will begin to protect themselves by reducing their water use, which can potentially reduce crop yields.


Hydrological Drought is associated with the effect of low rainfall on water levels in rivers, reservoirs, lakes, and aquifers. Hydrological droughts are usually noticed some time after meteorological droughts. First, precipitation decreases and after some time, water levels in rivers and lakes drop. Hydrological drought affects uses that depend on water levels. Changes in water levels affect ecosystems, hydroelectric power generation, and recreational, industrial and urban water use. A minor drought may affect small streams causing low streamflows or drying. A major drought could impact surface storage, lakes, and reservoirs thereby affecting water quality and causing municipal and agricultural water supply problems.

Rainfall also recharges groundwater aquifers through infiltration through the soil and run-off into streams and rivers. Once groundwater and surface waters are significantly impacted by lack of precipitation, a "hydrologic drought" occurs. Aquifer declines can range from a quick response (shallow sand) to impacts extending over multiple years. Impacts can include depletion of shallow depth wells, drying of farm dugouts, and changes to ground water quality.


Socioeconomic Drought occurs when the supply fails to meet the demand for an economic good(s) such as domestic water supplies, hay/forage, food grains, fish, and hydroelectric power, due to weather related water supply shortages from one or both of natural or managed water systems. At any time during meteorological, hydrological, or agricultural droughts, a socioeconomic drought can occur.

Attachments

1. Precipitation Indicator (Percent of 1, 3 and 12 month median precipitation)



THREE MONTH PRECIPITATION INDICATOR June 2014



Percent of Median

- 0-40 (Extreme)
- 41-60 (Severe)
- 61-80 (Moderate)
- > 80 (Normal)


0 25 50 100 150 200
Kilometers




Published by: Manitoba Conservation and Water Stewardship
 Projection: NAD 83 UTM Zone 14 N
 Printed: July 2014
 This map is for illustration purposes only

TWELVE MONTH PRECIPITATION INDICATOR

June 2014



2. Monthly Flow Indicator (lower 10th-20th-35th monthly flow percentile)





3. Water Supply Reservoir Status (Southern and Western)

	Water Supply Reservoir Levels and Storages							
	July 8, 2014							
Lake or Reservoir	Community	Target Level (feet)	Latest Observed Level (feet)	Observed date	Supply Status (Recent - Target) (feet)	Storage at Target Level (acre-feet)	Storage at Observed Level (acre-feet)	Supply Status (observed storage/target storage) (%)
Elgin	Elgin	1,532.00	1,532.04	March 24, 2014	0.04	520	523	101%
Goudney (Pilot Mound)	Pilot Mound	1,482.00	1,482.44	July 1, 2014	0.44	450	472	105%
Lake of the Prairies (Shellmouth)*	Brandon, Portage	1,402.50	1,415.66	July 7, 2014	13.16	300,000	514,083	171%
Manitou (Mary Jane)	Manitou	1,537.00	1,537.07	July 1, 2014	0.07	1,150	1,152	100%
Minnewasta (Morden)**	Morden	1,082.00	1,082.15	July 2, 2014	0.15	3,150	3,172	101%
Rapid City	Rapid City	1,573.50	1,573.73	May 2, 2014	0.23	200	216	108%
Lake Wahtopanah (Rivers)	Rivers	1,536.00	1,537.73	June 9, 2014	1.73	24,500	28,393	116%
Stephenfield	Carman	972.00	972.90	July 2, 2014	0.90	3,810	4,232	111%
Turtlehead (Deloraine)	Deloraine	1,772.00	1,775.16	July 1, 2014	3.16	1,400	1,652	118%
Vermilion	Dauphin	1,274.00	1,277.05	June 29, 2014	3.05	2,600	3,500	135%

* Summer Target level and storage. ** Corrected

4. Environment Canada 3 Month Outlook



5. Major River Basin

